

Hi!


Introducing SModelS - with an application to light neutralino DM

Suchita Kulkarni (LPSC, Grenoble)

based on:

- 1) work in progress with W. Waltenberger, U. Laa, A. Lessa, D. Proschofsky, S. Kraml
- 2) arxiv:1308.3735 [hep-ph] (accepted PLB) with G. Belanger, G. Drieu La Rochelle, B. Dumont, R. Godbole, S. Kraml

The scene



- Hunt for BSM physics is strong from the smallest to largest scales
- Many new and interesting results from astrophysics and collider searches exist and they must be taken into account to test a BSM theory
- Many BSM theories and no conclusive evidence for any of them
- The LHC searches are carried out in many channels and for many BSM models, here we focus on SUSY
- How does the scene for SUSY at LHC look like?

ATLAS SUSY Searches* - 95% CL Lower Limits

Status: SUSY 2013

ATLAS Preliminary

$$\int \mathcal{L} dt = (4.6 - 22.9) \text{ fb}^{-1} \quad \sqrt{s} = 7, 8 \text{ TeV}$$

Model	e, μ, τ, γ	Jets	E_T^{miss}	$\int \mathcal{L} dt [\text{fb}^{-1}]$	Mass limit	Reference		
Inclusive Searches	MSUGRA/CMSSM	0	2-6 jets	Yes	20.3	\tilde{q}, \tilde{g} 1.7 TeV	$m(\tilde{q})=m(\tilde{g})$	ATLAS-CONF-2013-047
	MSUGRA/CMSSM	1 e, μ	3-6 jets	Yes	20.3	\tilde{g} 1.2 TeV	any $m(\tilde{q})$	ATLAS-CONF-2013-062
	MSUGRA/CMSSM	0	7-10 jets	Yes	20.3	\tilde{g} 1.1 TeV	any $m(\tilde{q})$	1308.1841
	$\tilde{q}\tilde{q}, \tilde{q} \rightarrow q\tilde{\chi}_1^0$	0	2-6 jets	Yes	20.3	\tilde{q} 740 GeV	$m(\tilde{\chi}_1^0)=0 \text{ GeV}$	ATLAS-CONF-2013-047
	$\tilde{g}\tilde{g}, \tilde{g} \rightarrow q\tilde{q}\tilde{\chi}_1^0$	0	2-6 jets	Yes	20.3	\tilde{g} 1.3 TeV	$m(\tilde{\chi}_1^0)=0 \text{ GeV}$	ATLAS-CONF-2013-047
	$\tilde{g}\tilde{g}, \tilde{g} \rightarrow qq\tilde{\chi}_1^\pm \rightarrow qqW^\pm\tilde{\chi}_1^0$	1 e, μ	3-6 jets	Yes	20.3	\tilde{g} 1.18 TeV	$m(\tilde{\chi}_1^0) < 200 \text{ GeV}, m(\tilde{\chi}^\pm)=0.5(m(\tilde{\chi}_1^0)+m(\tilde{g}))$	ATLAS-CONF-2013-062
	$\tilde{g}\tilde{g}, \tilde{g} \rightarrow qq(\ell\ell/\ell\nu/\nu\nu)\tilde{\chi}_1^0$	2 e, μ	0-3 jets	-	20.3	\tilde{g} 1.12 TeV	$m(\tilde{\chi}_1^0)=0 \text{ GeV}$	ATLAS-CONF-2013-089
	GMSB ($\tilde{\ell}$ NLSP)	2 e, μ	2-4 jets	Yes	4.7	\tilde{g} 1.24 TeV	$\tan\beta < 15$	1208.4688
	GMSB ($\tilde{\ell}$ NLSP)	1-2 τ	0-2 jets	Yes	20.7	\tilde{g} 1.4 TeV	$\tan\beta > 18$	ATLAS-CONF-2013-026
	GGM (bino NLSP)	2 γ	-	Yes	4.8	\tilde{g} 1.07 TeV	$m(\tilde{\chi}_1^0) > 50 \text{ GeV}$	1209.0753
	GGM (wino NLSP)	1 $e, \mu + \gamma$	-	Yes	4.8	\tilde{g} 619 GeV	$m(\tilde{\chi}_1^0) > 50 \text{ GeV}$	ATLAS-CONF-2012-144
	GGM (higgsino-bino NLSP)	γ	1 b	Yes	4.8	\tilde{g} 900 GeV	$m(\tilde{\chi}_1^0) > 220 \text{ GeV}$	1211.1167
GGM (higgsino NLSP)	2 $e, \mu (Z)$	0-3 jets	Yes	5.8	\tilde{g} 690 GeV	$m(\tilde{H}) > 200 \text{ GeV}$	ATLAS-CONF-2012-152	
Gravitino LSP	0	mono-jet	Yes	10.5	$F^{1/2}$ scale 645 GeV	$m(\tilde{g}) > 10^{-4} \text{ eV}$	ATLAS-CONF-2012-147	
3 rd gen. \tilde{g} med.	$\tilde{g} \rightarrow b\tilde{b}\tilde{\chi}_1^0$	0	3 b	Yes	20.1	\tilde{g} 1.2 TeV	$m(\tilde{\chi}_1^0) < 600 \text{ GeV}$	ATLAS-CONF-2013-061
	$\tilde{g} \rightarrow t\tilde{t}\tilde{\chi}_1^0$	0	7-10 jets	Yes	20.3	\tilde{g} 1.1 TeV	$m(\tilde{\chi}_1^0) < 350 \text{ GeV}$	1308.1841
	$\tilde{g} \rightarrow t\tilde{t}\tilde{\chi}_1^\pm$	0-1 e, μ	3 b	Yes	20.1	\tilde{g} 1.34 TeV	$m(\tilde{\chi}_1^0) < 400 \text{ GeV}$	ATLAS-CONF-2013-061
	$\tilde{g} \rightarrow b\tilde{t}\tilde{\chi}_1^\pm$	0-1 e, μ	3 b	Yes	20.1	\tilde{g} 1.3 TeV	$m(\tilde{\chi}_1^0) < 300 \text{ GeV}$	ATLAS-CONF-2013-061
	3 rd gen. squarks direct production	$\tilde{b}_1\tilde{b}_1, \tilde{b}_1 \rightarrow b\tilde{\chi}_1^0$	0	2 b	Yes	20.1	\tilde{b}_1 100-620 GeV	$m(\tilde{\chi}_1^0) < 90 \text{ GeV}$
$\tilde{b}_1\tilde{b}_1, \tilde{b}_1 \rightarrow t\tilde{\chi}_1^\pm$		2 $e, \mu (SS)$	0-3 b	Yes	20.7	\tilde{b}_1 275-430 GeV	$m(\tilde{\chi}_1^\pm)=2 m(\tilde{\chi}_1^0)$	ATLAS-CONF-2013-007
$\tilde{t}_1\tilde{t}_1$ (light), $\tilde{t}_1 \rightarrow b\tilde{\chi}_1^\pm$		1-2 e, μ	1-2 b	Yes	4.7	\tilde{t}_1 110-167 GeV	$m(\tilde{\chi}_1^\pm)=55 \text{ GeV}$	1208.4305, 1209.2102
$\tilde{t}_1\tilde{t}_1$ (light), $\tilde{t}_1 \rightarrow Wb\tilde{\chi}_1^\pm$		2 e, μ	0-2 jets	Yes	20.3	\tilde{t}_1 130-220 GeV	$m(\tilde{\chi}_1^\pm) = m(\tilde{t}_1) - m(W) - 50 \text{ GeV}, m(\tilde{t}_1) < m(\tilde{\chi}_1^\pm)$	ATLAS-CONF-2013-048
$\tilde{t}_1\tilde{t}_1$ (medium), $\tilde{t}_1 \rightarrow t\tilde{\chi}_1^0$		2 e, μ	2 jets	Yes	20.3	\tilde{t}_1 225-525 GeV	$m(\tilde{\chi}_1^0)=0 \text{ GeV}$	ATLAS-CONF-2013-065
$\tilde{t}_1\tilde{t}_1$ (medium), $\tilde{t}_1 \rightarrow b\tilde{\chi}_1^\pm$		0	2 b	Yes	20.1	\tilde{t}_1 150-580 GeV	$m(\tilde{\chi}_1^0) < 200 \text{ GeV}, m(\tilde{\chi}_1^\pm) - m(\tilde{\chi}_1^0) = 5 \text{ GeV}$	1308.2631
$\tilde{t}_1\tilde{t}_1$ (heavy), $\tilde{t}_1 \rightarrow t\tilde{\chi}_1^0$		1 e, μ	1 b	Yes	20.7	\tilde{t}_1 200-610 GeV	$m(\tilde{\chi}_1^0)=0 \text{ GeV}$	ATLAS-CONF-2013-037
$\tilde{t}_1\tilde{t}_1$ (heavy), $\tilde{t}_1 \rightarrow t\tilde{\chi}_1^\pm$		0	2 b	Yes	20.5	\tilde{t}_1 320-660 GeV	$m(\tilde{\chi}_1^0)=0 \text{ GeV}$	ATLAS-CONF-2013-024
$\tilde{t}_1\tilde{t}_1, \tilde{t}_1 \rightarrow c\tilde{\chi}_1^0$		0	mono-jet/ c -tag	Yes	20.3	\tilde{t}_1 90-200 GeV	$m(\tilde{t}_1) - m(\tilde{\chi}_1^0) < 85 \text{ GeV}$	ATLAS-CONF-2013-068
$\tilde{t}_1\tilde{t}_1$ (natural GMSB)		2 $e, \mu (Z)$	1 b	Yes	20.7	\tilde{t}_1 500 GeV	$m(\tilde{\chi}_1^0) > 150 \text{ GeV}$	ATLAS-CONF-2013-025
$\tilde{t}_2\tilde{t}_2, \tilde{t}_2 \rightarrow \tilde{t}_1 + Z$		3 $e, \mu (Z)$	1 b	Yes	20.7	\tilde{t}_2 271-520 GeV	$m(\tilde{t}_1) = m(\tilde{\chi}_1^0) + 180 \text{ GeV}$	ATLAS-CONF-2013-025
EW direct	$\tilde{\ell}_{L,R}\tilde{\ell}_{L,R}, \tilde{\ell} \rightarrow \ell\tilde{\chi}_1^0$	2 e, μ	0	Yes	20.3	$\tilde{\ell}$ 85-315 GeV	$m(\tilde{\chi}_1^0)=0 \text{ GeV}$	ATLAS-CONF-2013-049
	$\tilde{\chi}_1^\pm\tilde{\chi}_1^\pm, \tilde{\chi}_1^\pm \rightarrow \tilde{\ell}\nu(\ell\bar{\nu})$	2 e, μ	0	Yes	20.3	$\tilde{\chi}_1^\pm$ 125-450 GeV	$m(\tilde{\chi}_1^0)=0 \text{ GeV}, m(\tilde{\ell}, \bar{\nu})=0.5(m(\tilde{\chi}_1^\pm)+m(\tilde{\chi}_1^0))$	ATLAS-CONF-2013-049
	$\tilde{\chi}_1^\pm\tilde{\chi}_1^\pm, \tilde{\chi}_1^\pm \rightarrow \tilde{\tau}\nu(\tau\bar{\nu})$	2 τ	-	Yes	20.7	$\tilde{\chi}_1^\pm$ 180-330 GeV	$m(\tilde{\chi}_1^0)=0 \text{ GeV}, m(\tilde{\tau}, \bar{\nu})=0.5(m(\tilde{\chi}_1^\pm)+m(\tilde{\chi}_1^0))$	ATLAS-CONF-2013-028
	$\tilde{\chi}_1^\pm\tilde{\chi}_2^0 \rightarrow \tilde{\ell}_L\nu\tilde{\ell}_L(\bar{\nu}\nu), \ell\bar{\nu}\tilde{\ell}_L(\ell\bar{\nu}\nu)$	3 e, μ	0	Yes	20.7	$\tilde{\chi}_1^\pm, \tilde{\chi}_2^0$ 600 GeV	$m(\tilde{\chi}_1^\pm)=m(\tilde{\chi}_2^0), m(\tilde{\chi}_1^0)=0, m(\tilde{\ell}, \bar{\nu})=0.5(m(\tilde{\chi}_1^\pm)+m(\tilde{\chi}_1^0))$	ATLAS-CONF-2013-035
	$\tilde{\chi}_1^\pm\tilde{\chi}_2^0 \rightarrow W\tilde{\chi}_1^0 Z\tilde{\chi}_1^0$	3 e, μ	0	Yes	20.7	$\tilde{\chi}_1^\pm, \tilde{\chi}_2^0$ 315 GeV	$m(\tilde{\chi}_1^\pm)=m(\tilde{\chi}_2^0), m(\tilde{\chi}_1^0)=0, \text{sleptons decoupled}$	ATLAS-CONF-2013-035
	$\tilde{\chi}_1^\pm\tilde{\chi}_2^0 \rightarrow W\tilde{\chi}_1^0 h\tilde{\chi}_1^0$	1 e, μ	2 b	Yes	20.3	$\tilde{\chi}_1^\pm, \tilde{\chi}_2^0$ 285 GeV	$m(\tilde{\chi}_1^\pm)=m(\tilde{\chi}_2^0), m(\tilde{\chi}_1^0)=0, \text{sleptons decoupled}$	ATLAS-CONF-2013-093
	Long-lived particles	Direct $\tilde{\chi}_1^\pm\tilde{\chi}_1^\pm$ prod., long-lived $\tilde{\chi}_1^\pm$	Disapp. trk	1 jet	Yes	20.3	$\tilde{\chi}_1^\pm$ 270 GeV	$m(\tilde{\chi}_1^\pm) - m(\tilde{\chi}_1^0) = 160 \text{ MeV}, \tau(\tilde{\chi}_1^\pm) = 0.2 \text{ ns}$
Stable, stopped \tilde{g} R-hadron		0	1-5 jets	Yes	22.9	\tilde{g} 832 GeV	$m(\tilde{\chi}_1^0)=100 \text{ GeV}, 10 \mu\text{s} < \tau(\tilde{g}) < 1000 \text{ s}$	ATLAS-CONF-2013-057
GMSB, stable $\tilde{\tau}, \tilde{\chi}_1^0 \rightarrow \tilde{\tau}(\tilde{e}, \tilde{\mu}) + \tau(e, \mu)$		1-2 μ	-	-	15.9	$\tilde{\chi}_1^0$ 475 GeV	$10 < \tan\beta < 50$	ATLAS-CONF-2013-058
GMSB, $\tilde{\chi}_1^0 \rightarrow \gamma\tilde{G}$, long-lived $\tilde{\chi}_1^0$		2 γ	-	Yes	4.7	$\tilde{\chi}_1^0$ 230 GeV	$0.4 < \tau(\tilde{\chi}_1^0) < 2 \text{ ns}$	1304.6310
$\tilde{q}\tilde{q}, \tilde{\chi}_1^0 \rightarrow qq\mu$ (RPV)		1 μ , displ. vtx	-	-	20.3	\tilde{q} 1.0 TeV	$1.5 < c\tau < 156 \text{ mm}, \text{BR}(\mu)=1, m(\tilde{\chi}_1^0)=108 \text{ GeV}$	ATLAS-CONF-2013-092
RPV	LFV $pp \rightarrow \tilde{\nu}_\tau + X, \tilde{\nu}_\tau \rightarrow e + \mu$	2 e, μ	-	-	4.6	$\tilde{\nu}_\tau$ 1.61 TeV	$\lambda'_{311}=0.10, \lambda_{132}=0.05$	1212.1272
	LFV $pp \rightarrow \tilde{\nu}_\tau + X, \tilde{\nu}_\tau \rightarrow e(\mu) + \tau$	1 $e, \mu + \tau$	-	-	4.6	$\tilde{\nu}_\tau$ 1.1 TeV	$\lambda'_{311}=0.10, \lambda_{1(2)33}=0.05$	1212.1272
	Bilinear RPV CMSSM	1 e, μ	7 jets	Yes	4.7	\tilde{q}, \tilde{g} 1.2 TeV	$m(\tilde{q})=m(\tilde{g}), c\tau_{LSP} < 1 \text{ mm}$	ATLAS-CONF-2012-140
	$\tilde{\chi}_1^\pm\tilde{\chi}_1^\pm, \tilde{\chi}_1^\pm \rightarrow W\tilde{\chi}_1^0, \tilde{\chi}_1^\pm \rightarrow ee\tilde{\nu}_\mu, e\mu\tilde{\nu}_e$	4 e, μ	-	Yes	20.7	$\tilde{\chi}_1^\pm$ 760 GeV	$m(\tilde{\chi}_1^0) > 300 \text{ GeV}, \lambda_{121} > 0$	ATLAS-CONF-2013-036
	$\tilde{\chi}_1^\pm\tilde{\chi}_1^\pm, \tilde{\chi}_1^\pm \rightarrow W\tilde{\chi}_1^0, \tilde{\chi}_1^\pm \rightarrow \tau\tau\tilde{\nu}_e, e\tau\tilde{\nu}_\tau$	3 $e, \mu + \tau$	-	Yes	20.7	$\tilde{\chi}_1^\pm$ 350 GeV	$m(\tilde{\chi}_1^0) > 80 \text{ GeV}, \lambda_{133} > 0$	ATLAS-CONF-2013-036
	$\tilde{g} \rightarrow qq\tilde{q}$	0	6-7 jets	-	20.3	\tilde{g} 916 GeV	$\text{BR}(t)=\text{BR}(b)=\text{BR}(c)=0\%$	ATLAS-CONF-2013-091
	$\tilde{g} \rightarrow \tilde{t}_1 t, \tilde{t}_1 \rightarrow bs$	2 $e, \mu (SS)$	0-3 b	Yes	20.7	\tilde{g} 880 GeV		ATLAS-CONF-2013-007
Other	Scalar gluon pair, $\text{sgluon} \rightarrow q\tilde{q}$	0	4 jets	-	4.6	sgluon 100-287 GeV	incl. limit from 1110.2693	1210.4826
	Scalar gluon pair, $\text{sgluon} \rightarrow t\tilde{t}$	2 $e, \mu (SS)$	1 b	Yes	14.3	sgluon 800 GeV		ATLAS-CONF-2013-051
	WIMP interaction (D5, Dirac χ)	0	mono-jet	Yes	10.5	M^* scale 704 GeV	$m(\chi) < 80 \text{ GeV}$, limit of $< 687 \text{ GeV}$ for D8	ATLAS-CONF-2012-147

$\sqrt{s} = 7 \text{ TeV}$ full data
 $\sqrt{s} = 8 \text{ TeV}$ partial data
 $\sqrt{s} = 8 \text{ TeV}$ full data

10⁻¹ 1 Mass scale [TeV]


*Only a selection of the available mass limits on new states or phenomena is shown. All limits quoted are observed minus 1 σ theoretical signal cross section uncertainty.

ATLAS SUSY Searches* - 95% CL Lower Limits

Status: SUSY 2013

ATLAS Preliminary

$$\int \mathcal{L} dt = (4.6 - 22.9) \text{ fb}^{-1} \quad \sqrt{s} = 7, 8 \text{ TeV}$$



$\sqrt{s} = 7$ TeV full data $\sqrt{s} = 8$ TeV partial data $\sqrt{s} = 8$ TeV full data

*Only a selection of the available mass limits on new states or phenomena is shown. All limits quoted are observed minus 1σ theoretical signal cross section uncertainty.

ATLAS SUSY Searches* - 95% CL Lower Limits

Status: SUSY 2013

ATLAS Preliminary

$$\int \mathcal{L} dt = (4.6 - 22.9) \text{ fb}^{-1} \quad \sqrt{s} = 7, 8 \text{ TeV}$$

Model	e, μ, τ, γ	Jets	E_T^{miss}	$\int \mathcal{L} dt [\text{fb}^{-1}]$	Mass limit	Reference		
Inclusive Searches	MSUGRA/CMSSM	0	2-6 jets	Yes	20.3	\tilde{q}, \tilde{g} 1.7 TeV	$m(\tilde{q})=m(\tilde{g})$	ATLAS-CONF-2013-047
	MSUGRA/CMSSM	1 e, μ	3-6 jets	Yes	20.3	\tilde{g} 1.2 TeV	any $m(\tilde{q})$	ATLAS-CONF-2013-062
	MSUGRA/CMSSM	0	7-10 jets	Yes	20.3	\tilde{g} 1.1 TeV	any $m(\tilde{q})$	1308.1841
	$\tilde{q}\tilde{q}, \tilde{q} \rightarrow q\tilde{\chi}_1^0$	0	2-6 jets	Yes	20.3	\tilde{q} 740 GeV	$m(\tilde{\chi}_1^0)=0 \text{ GeV}$	ATLAS-CONF-2013-047
	$\tilde{g}\tilde{g}, \tilde{g} \rightarrow q\tilde{q}\tilde{\chi}_1^0$	0	2-6 jets	Yes	20.3	\tilde{g} 1.3 TeV	$m(\tilde{\chi}_1^0)=0 \text{ GeV}$	ATLAS-CONF-2013-047
	$\tilde{g}\tilde{g}, \tilde{g} \rightarrow qq\tilde{\chi}_1^\pm \rightarrow qqW^\pm\tilde{\chi}_1^0$	1 e, μ	3-6 jets	Yes	20.3	\tilde{g} 1.18 TeV	$m(\tilde{\chi}_1^0) < 200 \text{ GeV}, m(\tilde{\chi}^\pm)=0.5(m(\tilde{\chi}_1^0)+m(\tilde{g}))$	ATLAS-CONF-2013-062
	$\tilde{g}\tilde{g}, \tilde{g} \rightarrow qq(\ell\ell/\ell\nu/\nu\nu)\tilde{\chi}_1^0$	2 e, μ	0-3 jets	-	20.3	\tilde{g} 1.12 TeV	$m(\tilde{\chi}_1^0)=0 \text{ GeV}$	ATLAS-CONF-2013-089
	GMSB ($\tilde{\ell}$ NLSP)	2 e, μ	2-4 jets	Yes	4.7	\tilde{g} 1.24 TeV	$\tan\beta < 15$	1208.4688
	GMSB ($\tilde{\ell}$ NLSP)	1-2 τ	0-2 jets	Yes	20.7	\tilde{g} 1.4 TeV	$\tan\beta > 18$	ATLAS-CONF-2013-026
	GGM (bino NLSP)	2 γ	-	Yes	4.8	\tilde{g} 1.07 TeV	$m(\tilde{\chi}_1^0) > 50 \text{ GeV}$	1209.0753
	GGM (wino NLSP)	1 $e, \mu + \gamma$	-	Yes	4.8	\tilde{g} 619 GeV	$m(\tilde{\chi}_1^0) > 50 \text{ GeV}$	ATLAS-CONF-2012-144
	GGM (higgsino-bino NLSP)	γ	1 b	Yes	4.8	\tilde{g} 900 GeV	$m(\tilde{\chi}_1^0) > 220 \text{ GeV}$	1211.1167
GGM (higgsino NLSP)	2 $e, \mu (Z)$	0-3 jets	Yes	5.8	\tilde{g} 690 GeV	$m(\tilde{H}) > 200 \text{ GeV}$	ATLAS-CONF-2012-152	
Gravitino LSP	0	mono-jet	Yes	10.5	$F^{1/2}$ scale 645 GeV	$m(\tilde{g}) > 10^{-4} \text{ eV}$	ATLAS-CONF-2012-147	
3 rd gen. \tilde{g} med.	$\tilde{g} \rightarrow b\tilde{b}\tilde{\chi}_1^0$	0	3 b	Yes	20.1	\tilde{g} 1.2 TeV	$m(\tilde{\chi}_1^0) < 600 \text{ GeV}$	ATLAS-CONF-2013-061
	$\tilde{g} \rightarrow t\tilde{t}\tilde{\chi}_1^0$	0	7-10 jets	Yes	20.3	\tilde{g} 1.1 TeV	$m(\tilde{\chi}_1^0) < 350 \text{ GeV}$	1308.1841
	$\tilde{g} \rightarrow t\tilde{t}\tilde{\chi}_1^\pm$	0-1 e, μ	3 b	Yes	20.1	\tilde{g} 1.34 TeV	$m(\tilde{\chi}_1^0) < 400 \text{ GeV}$	ATLAS-CONF-2013-061
	$\tilde{g} \rightarrow b\tilde{t}\tilde{\chi}_1^\pm$	0-1 e, μ	3 b	Yes	20.1	\tilde{g} 1.3 TeV	$m(\tilde{\chi}_1^0) < 300 \text{ GeV}$	ATLAS-CONF-2013-061
	3 rd gen. squarks direct production	$\tilde{b}_1\tilde{b}_1, \tilde{b}_1 \rightarrow b\tilde{\chi}_1^0$	0	2 b	Yes	20.1	\tilde{b}_1 100-620 GeV	$m(\tilde{\chi}_1^0) < 90 \text{ GeV}$
$\tilde{b}_1\tilde{b}_1, \tilde{b}_1 \rightarrow t\tilde{\chi}_1^\pm$		2 e, μ (SS)	0-3 b	Yes	20.7	\tilde{b}_1 275-430 GeV	$m(\tilde{\chi}_1^\pm)=2 m(\tilde{\chi}_1^0)$	ATLAS-CONF-2013-007
$\tilde{t}_1\tilde{t}_1$ (light), $\tilde{t}_1 \rightarrow b\tilde{\chi}_1^\pm$		1-2 e, μ	1-2 b	Yes	4.7	\tilde{t}_1 110-167 GeV	$m(\tilde{\chi}_1^\pm)=55 \text{ GeV}$	1208.4305, 1209.2102
$\tilde{t}_1\tilde{t}_1$ (light), $\tilde{t}_1 \rightarrow Wb\tilde{\chi}_1^\pm$		2 e, μ	0-2 jets	Yes	20.3	\tilde{t}_1 130-220 GeV	$m(\tilde{\chi}_1^\pm) = m(\tilde{t}_1) - m(W) - 50 \text{ GeV}, m(\tilde{t}_1) < m(\tilde{\chi}_1^\pm)$	ATLAS-CONF-2013-048
$\tilde{t}_1\tilde{t}_1$ (medium), $\tilde{t}_1 \rightarrow t\tilde{\chi}_1^0$		2 e, μ	2 jets	Yes	20.3	\tilde{t}_1 225-525 GeV	$m(\tilde{\chi}_1^0)=0 \text{ GeV}$	ATLAS-CONF-2013-065
$\tilde{t}_1\tilde{t}_1$ (medium), $\tilde{t}_1 \rightarrow b\tilde{\chi}_1^\pm$		0	2 b	Yes	20.1	\tilde{t}_1 150-580 GeV	$m(\tilde{\chi}_1^0) < 200 \text{ GeV}, m(\tilde{\chi}_1^\pm) - m(\tilde{\chi}_1^0) = 5 \text{ GeV}$	1308.2631
$\tilde{t}_1\tilde{t}_1$ (heavy), $\tilde{t}_1 \rightarrow t\tilde{\chi}_1^0$		1 e, μ	1 b	Yes	20.7	\tilde{t}_1 200-610 GeV	$m(\tilde{\chi}_1^0)=0 \text{ GeV}$	ATLAS-CONF-2013-037
$\tilde{t}_1\tilde{t}_1$ (heavy), $\tilde{t}_1 \rightarrow t\tilde{\chi}_1^\pm$		0	2 b	Yes	20.5	\tilde{t}_1 320-660 GeV	$m(\tilde{\chi}_1^0)=0 \text{ GeV}$	ATLAS-CONF-2013-024
$\tilde{t}_1\tilde{t}_1, \tilde{t}_1 \rightarrow c\tilde{\chi}_1^0$		0	mono-jet/ c -tag	Yes	20.3	\tilde{t}_1 90-200 GeV	$m(\tilde{t}_1) - m(\tilde{\chi}_1^0) < 85 \text{ GeV}$	ATLAS-CONF-2013-068
$\tilde{t}_1\tilde{t}_1$ (natural GMSB)		2 $e, \mu (Z)$	1 b	Yes	20.7	\tilde{t}_1 500 GeV	$m(\tilde{\chi}_1^0) > 150 \text{ GeV}$	ATLAS-CONF-2013-025
$\tilde{t}_2\tilde{t}_2, \tilde{t}_2 \rightarrow \tilde{t}_1 + Z$		3 $e, \mu (Z)$	1 b	Yes	20.7	\tilde{t}_2 271-520 GeV	$m(\tilde{t}_1) = m(\tilde{\chi}_1^0) + 180 \text{ GeV}$	ATLAS-CONF-2013-025
EW direct	$\tilde{\ell}_{L,R}\tilde{\ell}_{L,R}, \tilde{\ell} \rightarrow \ell\tilde{\chi}_1^0$	2 e, μ	0	Yes	20.3	$\tilde{\ell}$ 85-315 GeV	$m(\tilde{\chi}_1^0)=0 \text{ GeV}$	ATLAS-CONF-2013-049
	$\tilde{\chi}_1^\pm\tilde{\chi}_1^\pm, \tilde{\chi}_1^\pm \rightarrow \tilde{\ell}\nu(\ell\bar{\nu})$	2 e, μ	0	Yes	20.3	$\tilde{\chi}_1^\pm$ 125-450 GeV	$m(\tilde{\chi}_1^0)=0 \text{ GeV}, m(\tilde{\ell}, \bar{\nu})=0.5(m(\tilde{\chi}_1^\pm)+m(\tilde{\chi}_1^0))$	ATLAS-CONF-2013-049
	$\tilde{\chi}_1^\pm\tilde{\chi}_1^\pm, \tilde{\chi}_1^\pm \rightarrow \tilde{\tau}\nu(\tau\bar{\nu})$	2 τ	-	Yes	20.7	$\tilde{\chi}_1^\pm$ 180-330 GeV	$m(\tilde{\chi}_1^0)=0 \text{ GeV}, m(\tilde{\tau}, \bar{\nu})=0.5(m(\tilde{\chi}_1^\pm)+m(\tilde{\chi}_1^0))$	ATLAS-CONF-2013-028
	$\tilde{\chi}_1^\pm\tilde{\chi}_2^0 \rightarrow \tilde{\ell}_L\nu\tilde{\ell}_L(\bar{\nu}\nu), \ell\bar{\nu}\tilde{\ell}_L(\bar{\nu}\nu)$	3 e, μ	0	Yes	20.7	$\tilde{\chi}_1^\pm, \tilde{\chi}_2^0$ 600 GeV	$m(\tilde{\chi}_1^\pm)=m(\tilde{\chi}_2^0), m(\tilde{\chi}_1^0)=0, m(\tilde{\ell}, \bar{\nu})=0.5(m(\tilde{\chi}_1^\pm)+m(\tilde{\chi}_1^0))$	ATLAS-CONF-2013-035
	$\tilde{\chi}_1^\pm\tilde{\chi}_2^0 \rightarrow W\tilde{\chi}_1^0 Z\tilde{\chi}_1^0$	3 e, μ	0	Yes	20.7	$\tilde{\chi}_1^\pm, \tilde{\chi}_2^0$ 315 GeV	$m(\tilde{\chi}_1^\pm)=m(\tilde{\chi}_2^0), m(\tilde{\chi}_1^0)=0, \text{sleptons decoupled}$	ATLAS-CONF-2013-035
	$\tilde{\chi}_1^\pm\tilde{\chi}_2^0 \rightarrow W\tilde{\chi}_1^0 h\tilde{\chi}_1^0$	1 e, μ	2 b	Yes	20.3	$\tilde{\chi}_1^\pm, \tilde{\chi}_2^0$ 285 GeV	$m(\tilde{\chi}_1^\pm)=m(\tilde{\chi}_2^0), m(\tilde{\chi}_1^0)=0, \text{sleptons decoupled}$	ATLAS-CONF-2013-093
	Long-lived particles	Direct $\tilde{\chi}_1^\pm\tilde{\chi}_1^\pm$ prod., long-lived $\tilde{\chi}_1^\pm$	Disapp. trk	1 jet	Yes	20.3	$\tilde{\chi}_1^\pm$ 270 GeV	$m(\tilde{\chi}_1^\pm) - m(\tilde{\chi}_1^0) = 160 \text{ MeV}, \tau(\tilde{\chi}_1^\pm) = 0.2 \text{ ns}$
Stable, stopped \tilde{g} R-hadron		0	1-5 jets	Yes	22.9	\tilde{g} 832 GeV	$m(\tilde{\chi}_1^0)=100 \text{ GeV}, 10 \mu\text{s} < \tau(\tilde{g}) < 1000 \text{ s}$	ATLAS-CONF-2013-057
GMSB, stable $\tilde{\tau}, \tilde{\chi}_1^0 \rightarrow \tilde{\tau}(\tilde{e}, \tilde{\mu}) + \tau(e, \mu)$		1-2 μ	-	-	15.9	$\tilde{\chi}_1^0$ 475 GeV	$10 < \tan\beta < 50$	ATLAS-CONF-2013-058
GMSB, $\tilde{\chi}_1^0 \rightarrow \gamma\tilde{G}$, long-lived $\tilde{\chi}_1^0$		2 γ	-	Yes	4.7	$\tilde{\chi}_1^0$ 230 GeV	$0.4 < \tau(\tilde{\chi}_1^0) < 2 \text{ ns}$	1304.6310
$\tilde{q}\tilde{q}, \tilde{\chi}_1^0 \rightarrow qq\mu$ (RPV)		1 μ , displ. vtx	-	-	20.3	\tilde{q} 1.0 TeV	$1.5 < c\tau < 156 \text{ mm}, \text{BR}(\mu)=1, m(\tilde{\chi}_1^0)=108 \text{ GeV}$	ATLAS-CONF-2013-092
RPV	LFV $pp \rightarrow \tilde{\nu}_\tau + X, \tilde{\nu}_\tau \rightarrow e + \mu$	2 e, μ	-	-	4.6	$\tilde{\nu}_\tau$ 1.61 TeV	$\lambda'_{311}=0.10, \lambda_{132}=0.05$	1212.1272
	LFV $pp \rightarrow \tilde{\nu}_\tau + X, \tilde{\nu}_\tau \rightarrow e(\mu) + \tau$	1 $e, \mu + \tau$	-	-	4.6	$\tilde{\nu}_\tau$ 1.1 TeV	$\lambda'_{311}=0.10, \lambda_{1(2)33}=0.05$	1212.1272
	Bilinear RPV CMSSM	1 e, μ	7 jets	Yes	4.7	\tilde{q}, \tilde{g} 1.2 TeV	$m(\tilde{q})=m(\tilde{g}), c\tau_{\text{LSP}} < 1 \text{ mm}$	ATLAS-CONF-2012-140
	$\tilde{\chi}_1^\pm\tilde{\chi}_1^\pm, \tilde{\chi}_1^\pm \rightarrow W\tilde{\chi}_1^0, \tilde{\chi}_1^\pm \rightarrow ee\tilde{\nu}_\mu, e\mu\tilde{\nu}_e$	4 e, μ	-	Yes	20.7	$\tilde{\chi}_1^\pm$ 760 GeV	$m(\tilde{\chi}_1^0) > 300 \text{ GeV}, \lambda_{121} > 0$	ATLAS-CONF-2013-036
	$\tilde{\chi}_1^\pm\tilde{\chi}_1^\pm, \tilde{\chi}_1^\pm \rightarrow W\tilde{\chi}_1^0, \tilde{\chi}_1^\pm \rightarrow \tau\tau\tilde{\nu}_e, e\tau\tilde{\nu}_\tau$	3 $e, \mu + \tau$	-	Yes	20.7	$\tilde{\chi}_1^\pm$ 350 GeV	$m(\tilde{\chi}_1^0) > 80 \text{ GeV}, \lambda_{133} > 0$	ATLAS-CONF-2013-036
	$\tilde{g} \rightarrow qq\tilde{q}$	0	6-7 jets	-	20.3	\tilde{g} 916 GeV	$\text{BR}(t)=\text{BR}(b)=\text{BR}(c)=0\%$	ATLAS-CONF-2013-091
	$\tilde{g} \rightarrow \tilde{t}_1 t, \tilde{t}_1 \rightarrow bs$	2 e, μ (SS)	0-3 b	Yes	20.7	\tilde{g} 880 GeV		ATLAS-CONF-2013-007
Other	Scalar gluon pair, $\text{sgluon} \rightarrow q\tilde{q}$	0	4 jets	-	4.6	sgluon 100-287 GeV	incl. limit from 1110.2693	1210.4826
	Scalar gluon pair, $\text{sgluon} \rightarrow t\tilde{t}$	2 e, μ (SS)	1 b	Yes	14.3	sgluon 800 GeV		ATLAS-CONF-2013-051
	WIMP interaction (D5, Dirac χ)	0	mono-jet	Yes	10.5	M^* scale 704 GeV	$m(\chi) < 80 \text{ GeV}, \text{limit of } < 687 \text{ GeV for D8}$	ATLAS-CONF-2012-147

$\sqrt{s} = 7 \text{ TeV}$ full data
 $\sqrt{s} = 8 \text{ TeV}$ partial data
 $\sqrt{s} = 8 \text{ TeV}$ full data

10⁻¹ 1 Mass scale [TeV]


*Only a selection of the available mass limits on new states or phenomena is shown. All limits quoted are observed minus 1 σ theoretical signal cross section uncertainty.

ATLAS SUSY Searches* - 95% CL Lower Limits

Status: SUSY 2013

ATLAS Preliminary

$$\int \mathcal{L} dt = (4.6 - 22.9) \text{ fb}^{-1} \quad \sqrt{s} = 7, 8 \text{ TeV}$$



200-300 GeV

$\sqrt{s} = 7 \text{ TeV}$ full data
 $\sqrt{s} = 8 \text{ TeV}$ partial data
 $\sqrt{s} = 8 \text{ TeV}$ full data

*Only a selection of the available mass limits on new states or phenomena is shown. All limits quoted are observed minus 1σ theoretical signal cross section uncertainty.

Behind the scene

Behind the scene

- Results based on several assumptions and are the best limits -- should be taken with a grain of salt for a generic model

ATLAS SUSY Searches* - 95% CL Lower Limits

Status: SUSY 2013

ATLAS Preliminary

$$\int \mathcal{L} dt = (4.6 - 22.9) \text{ fb}^{-1} \quad \sqrt{s} = 7, 8 \text{ TeV}$$

Model	e, μ, τ, γ	Jets	E_T^{miss}	$\int \mathcal{L} dt [\text{fb}^{-1}]$	Mass limit	Reference		
Inclusive Searches	MSUGRA/CMSSM	0	2-6 jets	Yes	20.3	\tilde{q}, \tilde{g} 1.7 TeV	$m(\tilde{q})=m(\tilde{g})$	ATLAS-CONF-2013-047
	MSUGRA/CMSSM	1 e, μ	3-6 jets	Yes	20.3	\tilde{g} 1.2 TeV	any $m(\tilde{q})$	ATLAS-CONF-2013-062
	MSUGRA/CMSSM	0	7-10 jets	Yes	20.3	\tilde{g} 1.1 TeV	any $m(\tilde{q})$	1308.1841
	$\tilde{q}\tilde{q}, \tilde{q} \rightarrow q\tilde{\chi}_1^0$	0	2-6 jets	Yes	20.3	\tilde{q} 740 GeV	$m(\tilde{\chi}_1^0)=0 \text{ GeV}$	ATLAS-CONF-2013-047
	$\tilde{g}\tilde{g}, \tilde{g} \rightarrow q\tilde{q}\tilde{\chi}_1^0$	0	2-6 jets	Yes	20.3	\tilde{g} 1.3 TeV	$m(\tilde{\chi}_1^0)=0 \text{ GeV}$	ATLAS-CONF-2013-047
	$\tilde{g}\tilde{g}, \tilde{g} \rightarrow qq\tilde{\chi}_1^\pm \rightarrow qqW^\pm\tilde{\chi}_1^0$	1 e, μ	3-6 jets	Yes	20.3	\tilde{g} 1.18 TeV	$m(\tilde{\chi}_1^0) < 200 \text{ GeV}, m(\tilde{\chi}^\pm)=0.5(m(\tilde{\chi}_1^0)+m(\tilde{g}))$	ATLAS-CONF-2013-062
	$\tilde{g}\tilde{g}, \tilde{g} \rightarrow qq(\ell\ell/\ell\nu/\nu\nu)\tilde{\chi}_1^0$	2 e, μ	0-3 jets	-	20.3	\tilde{g} 1.12 TeV	$m(\tilde{\chi}_1^0)=0 \text{ GeV}$	ATLAS-CONF-2013-089
	GMSB ($\tilde{\ell}$ NLSP)	2 e, μ	2-4 jets	Yes	4.7	\tilde{g} 1.24 TeV	$\tan\beta < 15$	1208.4688
	GMSB ($\tilde{\ell}$ NLSP)	1-2 τ	0-2 jets	Yes	20.7	\tilde{g} 1.4 TeV	$\tan\beta > 18$	ATLAS-CONF-2013-026
	GGM (bino NLSP)	2 γ	-	Yes	4.8	\tilde{g} 1.07 TeV	$m(\tilde{\chi}_1^0) > 50 \text{ GeV}$	1209.0753
	GGM (wino NLSP)	1 $e, \mu + \gamma$	-	Yes	4.8	\tilde{g} 619 GeV	$m(\tilde{\chi}_1^0) > 50 \text{ GeV}$	ATLAS-CONF-2012-144
	GGM (higgsino-bino NLSP)	γ	1 b	Yes	4.8	\tilde{g} 900 GeV	$m(\tilde{\chi}_1^0) > 220 \text{ GeV}$	1211.1167
GGM (higgsino NLSP)	2 $e, \mu (Z)$	0-3 jets	Yes	5.8	\tilde{g} 690 GeV	$m(\tilde{H}) > 200 \text{ GeV}$	ATLAS-CONF-2012-152	
Gravitino LSP	0	mono-jet	Yes	10.5	$F^{1/2}$ scale 645 GeV	$m(\tilde{g}) > 10^{-4} \text{ eV}$	ATLAS-CONF-2012-147	
3 rd gen. \tilde{g} med.	$\tilde{g} \rightarrow b\tilde{b}\tilde{\chi}_1^0$	0	3 b	Yes	20.1	\tilde{g} 1.2 TeV	$m(\tilde{\chi}_1^0) < 600 \text{ GeV}$	ATLAS-CONF-2013-061
	$\tilde{g} \rightarrow t\tilde{t}\tilde{\chi}_1^0$	0	7-10 jets	Yes	20.3	\tilde{g} 1.1 TeV	$m(\tilde{\chi}_1^0) < 350 \text{ GeV}$	1308.1841
	$\tilde{g} \rightarrow t\tilde{t}\tilde{\chi}_1^\pm$	0-1 e, μ	3 b	Yes	20.1	\tilde{g} 1.34 TeV	$m(\tilde{\chi}_1^0) < 400 \text{ GeV}$	ATLAS-CONF-2013-061
	$\tilde{g} \rightarrow b\tilde{t}\tilde{\chi}_1^\pm$	0-1 e, μ	3 b	Yes	20.1	\tilde{g} 1.3 TeV	$m(\tilde{\chi}_1^0) < 300 \text{ GeV}$	ATLAS-CONF-2013-061
	3 rd gen. squarks direct production	$\tilde{b}_1\tilde{b}_1, \tilde{b}_1 \rightarrow b\tilde{\chi}_1^0$	0	2 b	Yes	20.1	\tilde{b}_1 100-620 GeV	$m(\tilde{\chi}_1^0) < 90 \text{ GeV}$
$\tilde{b}_1\tilde{b}_1, \tilde{b}_1 \rightarrow t\tilde{\chi}_1^\pm$		2 e, μ (SS)	0-3 b	Yes	20.7	\tilde{b}_1 275-430 GeV	$m(\tilde{\chi}_1^\pm)=2 m(\tilde{\chi}_1^0)$	ATLAS-CONF-2013-007
$\tilde{t}_1\tilde{t}_1$ (light), $\tilde{t}_1 \rightarrow b\tilde{\chi}_1^\pm$		1-2 e, μ	1-2 b	Yes	4.7	\tilde{t}_1 110-167 GeV	$m(\tilde{\chi}_1^\pm)=55 \text{ GeV}$	1208.4305, 1209.2102
$\tilde{t}_1\tilde{t}_1$ (light), $\tilde{t}_1 \rightarrow Wb\tilde{\chi}_1^\pm$		2 e, μ	0-2 jets	Yes	20.3	\tilde{t}_1 130-220 GeV	$m(\tilde{\chi}_1^\pm) = m(\tilde{t}_1) - m(W) - 50 \text{ GeV}, m(\tilde{t}_1) < m(\tilde{\chi}_1^\pm)$	ATLAS-CONF-2013-048
$\tilde{t}_1\tilde{t}_1$ (medium), $\tilde{t}_1 \rightarrow t\tilde{\chi}_1^0$		2 e, μ	2 jets	Yes	20.3	\tilde{t}_1 225-525 GeV	$m(\tilde{\chi}_1^0)=0 \text{ GeV}$	ATLAS-CONF-2013-065
$\tilde{t}_1\tilde{t}_1$ (medium), $\tilde{t}_1 \rightarrow b\tilde{\chi}_1^\pm$		0	2 b	Yes	20.1	\tilde{t}_1 150-580 GeV	$m(\tilde{\chi}_1^0) < 200 \text{ GeV}, m(\tilde{\chi}_1^\pm) - m(\tilde{\chi}_1^0) = 5 \text{ GeV}$	1308.2631
$\tilde{t}_1\tilde{t}_1$ (heavy), $\tilde{t}_1 \rightarrow t\tilde{\chi}_1^0$		1 e, μ	1 b	Yes	20.7	\tilde{t}_1 200-610 GeV	$m(\tilde{\chi}_1^0)=0 \text{ GeV}$	ATLAS-CONF-2013-037
$\tilde{t}_1\tilde{t}_1$ (heavy), $\tilde{t}_1 \rightarrow t\tilde{\chi}_1^\pm$		0	2 b	Yes	20.5	\tilde{t}_1 320-660 GeV	$m(\tilde{\chi}_1^0)=0 \text{ GeV}$	ATLAS-CONF-2013-024
$\tilde{t}_1\tilde{t}_1, \tilde{t}_1 \rightarrow c\tilde{\chi}_1^0$		0	mono-jet/ c -tag	Yes	20.3	\tilde{t}_1 90-200 GeV	$m(\tilde{t}_1) - m(\tilde{\chi}_1^0) < 85 \text{ GeV}$	ATLAS-CONF-2013-068
$\tilde{t}_1\tilde{t}_1$ (natural GMSB)		2 $e, \mu (Z)$	1 b	Yes	20.7	\tilde{t}_1 500 GeV	$m(\tilde{\chi}_1^0) > 150 \text{ GeV}$	ATLAS-CONF-2013-025
$\tilde{t}_2\tilde{t}_2, \tilde{t}_2 \rightarrow \tilde{t}_1 + Z$		3 $e, \mu (Z)$	1 b	Yes	20.7	\tilde{t}_2 271-520 GeV	$m(\tilde{t}_1) = m(\tilde{\chi}_1^0) + 180 \text{ GeV}$	ATLAS-CONF-2013-025
EW direct	$\tilde{\ell}_{L,R}\tilde{\ell}_{L,R}, \tilde{\ell} \rightarrow \ell\tilde{\chi}_1^0$	2 e, μ	0	Yes	20.3	$\tilde{\ell}$ 85-315 GeV	$m(\tilde{\chi}_1^0)=0 \text{ GeV}$	ATLAS-CONF-2013-049
	$\tilde{\chi}_1^\pm\tilde{\chi}_1^\pm, \tilde{\chi}_1^\pm \rightarrow \tilde{\ell}\nu(\ell\bar{\nu})$	2 e, μ	0	Yes	20.3	$\tilde{\chi}_1^\pm$ 125-450 GeV	$m(\tilde{\chi}_1^0)=0 \text{ GeV}, m(\tilde{\ell}, \bar{\nu})=0.5(m(\tilde{\chi}_1^\pm)+m(\tilde{\chi}_1^0))$	ATLAS-CONF-2013-049
	$\tilde{\chi}_1^\pm\tilde{\chi}_1^\pm, \tilde{\chi}_1^\pm \rightarrow \tilde{\tau}\nu(\tau\bar{\nu})$	2 τ	-	Yes	20.7	$\tilde{\chi}_1^\pm$ 180-330 GeV	$m(\tilde{\chi}_1^0)=0 \text{ GeV}, m(\tilde{\tau}, \bar{\nu})=0.5(m(\tilde{\chi}_1^\pm)+m(\tilde{\chi}_1^0))$	ATLAS-CONF-2013-028
	$\tilde{\chi}_1^\pm\tilde{\chi}_2^0 \rightarrow \tilde{\ell}_L\nu\tilde{\ell}_L(\bar{\nu}\nu), \ell\bar{\nu}\tilde{\ell}_L(\bar{\nu}\nu)$	3 e, μ	0	Yes	20.7	$\tilde{\chi}_1^\pm, \tilde{\chi}_2^0$ 600 GeV	$m(\tilde{\chi}_1^\pm)=m(\tilde{\chi}_2^0), m(\tilde{\chi}_1^0)=0, m(\tilde{\ell}, \bar{\nu})=0.5(m(\tilde{\chi}_1^\pm)+m(\tilde{\chi}_1^0))$	ATLAS-CONF-2013-035
	$\tilde{\chi}_1^\pm\tilde{\chi}_2^0 \rightarrow W\tilde{\chi}_1^0 Z\tilde{\chi}_1^0$	3 e, μ	0	Yes	20.7	$\tilde{\chi}_1^\pm, \tilde{\chi}_2^0$ 315 GeV	$m(\tilde{\chi}_1^\pm)=m(\tilde{\chi}_2^0), m(\tilde{\chi}_1^0)=0, \text{sleptons decoupled}$	ATLAS-CONF-2013-035
	$\tilde{\chi}_1^\pm\tilde{\chi}_2^0 \rightarrow W\tilde{\chi}_1^0 h\tilde{\chi}_1^0$	1 e, μ	2 b	Yes	20.3	$\tilde{\chi}_1^\pm, \tilde{\chi}_2^0$ 285 GeV	$m(\tilde{\chi}_1^\pm)=m(\tilde{\chi}_2^0), m(\tilde{\chi}_1^0)=0, \text{sleptons decoupled}$	ATLAS-CONF-2013-093
Long-lived particles	Direct $\tilde{\chi}_1^\pm\tilde{\chi}_1^\pm$ prod., long-lived $\tilde{\chi}_1^\pm$	Disapp. trk	1 jet	Yes	20.3	$\tilde{\chi}_1^\pm$ 270 GeV	$m(\tilde{\chi}_1^\pm) - m(\tilde{\chi}_1^0) = 160 \text{ MeV}, \tau(\tilde{\chi}_1^\pm) = 0.2 \text{ ns}$	ATLAS-CONF-2013-069
	Stable, stopped \tilde{g} R-hadron	0	1-5 jets	Yes	22.9	\tilde{g} 832 GeV	$m(\tilde{\chi}_1^0)=100 \text{ GeV}, 10 \mu\text{s} < \tau(\tilde{g}) < 1000 \text{ s}$	ATLAS-CONF-2013-057
	GMSB, stable $\tilde{\tau}, \tilde{\chi}_1^0 \rightarrow \tilde{\tau}(\tilde{e}, \tilde{\mu}) + \tau(e, \mu)$	1-2 μ	-	-	15.9	$\tilde{\chi}_1^0$ 475 GeV	$10 < \tan\beta < 50$	ATLAS-CONF-2013-058
	GMSB, $\tilde{\chi}_1^0 \rightarrow \gamma\tilde{G}$, long-lived $\tilde{\chi}_1^0$	2 γ	-	Yes	4.7	$\tilde{\chi}_1^0$ 230 GeV	$0.4 < \tau(\tilde{\chi}_1^0) < 2 \text{ ns}$	1304.6310
$\tilde{q}\tilde{q}, \tilde{\chi}_1^0 \rightarrow qq\mu$ (RPV)	1 μ , displ. vtx	-	-	20.3	\tilde{q} 1.0 TeV	$1.5 < c\tau < 156 \text{ mm}, \text{BR}(\mu)=1, m(\tilde{\chi}_1^0)=108 \text{ GeV}$	ATLAS-CONF-2013-092	
RPV	LFV $pp \rightarrow \tilde{\nu}_\tau + X, \tilde{\nu}_\tau \rightarrow e + \mu$	2 e, μ	-	-	4.6	$\tilde{\nu}_\tau$ 1.61 TeV	$\lambda'_{311}=0.10, \lambda_{132}=0.05$	1212.1272
	LFV $pp \rightarrow \tilde{\nu}_\tau + X, \tilde{\nu}_\tau \rightarrow e(\mu) + \tau$	1 $e, \mu + \tau$	-	-	4.6	$\tilde{\nu}_\tau$ 1.1 TeV	$\lambda'_{311}=0.10, \lambda_{1(2)33}=0.05$	1212.1272
	Bilinear RPV CMSSM	1 e, μ	7 jets	Yes	4.7	\tilde{q}, \tilde{g} 1.2 TeV	$m(\tilde{q})=m(\tilde{g}), c\tau_{\text{LSP}} < 1 \text{ mm}$	ATLAS-CONF-2012-140
	$\tilde{\chi}_1^\pm\tilde{\chi}_1^\pm, \tilde{\chi}_1^\pm \rightarrow W\tilde{\chi}_1^0, \tilde{\chi}_1^\pm \rightarrow ee\tilde{\nu}_\mu, e\mu\tilde{\nu}_e$	4 e, μ	-	Yes	20.7	$\tilde{\chi}_1^\pm$ 760 GeV	$m(\tilde{\chi}_1^0) > 300 \text{ GeV}, \lambda_{121} > 0$	ATLAS-CONF-2013-036
	$\tilde{\chi}_1^\pm\tilde{\chi}_1^\pm, \tilde{\chi}_1^\pm \rightarrow W\tilde{\chi}_1^0, \tilde{\chi}_1^\pm \rightarrow \tau\tau\tilde{\nu}_e, e\tau\tilde{\nu}_\tau$	3 $e, \mu + \tau$	-	Yes	20.7	$\tilde{\chi}_1^\pm$ 350 GeV	$m(\tilde{\chi}_1^0) > 80 \text{ GeV}, \lambda_{133} > 0$	ATLAS-CONF-2013-036
	$\tilde{g} \rightarrow qq\tilde{q}$	0	6-7 jets	-	20.3	\tilde{g} 916 GeV	$\text{BR}(t)=\text{BR}(b)=\text{BR}(c)=0\%$	ATLAS-CONF-2013-091
	$\tilde{g} \rightarrow \tilde{t}_1 t, \tilde{t}_1 \rightarrow bs$	2 e, μ (SS)	0-3 b	Yes	20.7	\tilde{g} 880 GeV		ATLAS-CONF-2013-007
Other	Scalar gluon pair, $\text{sgluon} \rightarrow q\tilde{q}$	0	4 jets	-	4.6	sgluon 100-287 GeV	incl. limit from 1110.2693	1210.4826
	Scalar gluon pair, $\text{sgluon} \rightarrow t\tilde{t}$	2 e, μ (SS)	1 b	Yes	14.3	sgluon 800 GeV		ATLAS-CONF-2013-051
	WIMP interaction (D5, Dirac χ)	0	mono-jet	Yes	10.5	M^* scale 704 GeV	$m(\chi) < 80 \text{ GeV}, \text{limit of } < 687 \text{ GeV for D8}$	ATLAS-CONF-2012-147

$\sqrt{s} = 7 \text{ TeV}$ full data
 $\sqrt{s} = 8 \text{ TeV}$ partial data
 $\sqrt{s} = 8 \text{ TeV}$ full data


10⁻¹ 1 Mass scale [TeV]

*Only a selection of the available mass limits on new states or phenomena is shown. All limits quoted are observed minus 1 σ theoretical signal cross section uncertainty.

Behind the scene

- Results based on several assumptions and are the best limits -- should be taken with a grain of salt for a generic model
- Simplified Model Spectra (SMS) are a good way to represent a more generic picture
- The alternative to SMS is to re-implement the relevant analyses to test the SUSY scenario - typically time consuming, demands huge computing power but more accurate
- We take the simpler approach and stick to SMS results here

How to read a SMS result



ATLAS-CONF-2013-024

- 95% CS UL is the unfolded maximum amount of cross-section allowed for a specific decay chain and a mass combination

Can we have a centralized database of all the SMS results to check a given SUSY point in parameter space by decomposing it into SMS topologies?


Central concept of

SM_oodels

The logo for SM_oodels features the text "SMoodels" in a grey, rounded font. The letter "o" is replaced by a stylized orange particle symbol, specifically a gluon loop, which consists of a central circle with two wavy lines extending downwards from its bottom edge.

SModelS framework


- It assumes, for most experimental searches, the BSM model can be approximated by a sum over effective simplified models



- Current implementation assumes R-parity is conserved

SModelS framework

- Consider:




- The framework does not depend on characteristics of SUSY particles, can also be applied to decompose any BSM spectra of arbitrary complexity


A real life application

Do LHC results on the SUSY particles, Higgs signal strengths and constraints on DM from direct and indirect detection experiments rule out light neutralino DM?

Already many studies exist in literature, I'll not list them here



126 GeV Higgs at the LHC




- Four production modes ggF, VH, VBF, and ttH
- Five final states: $\gamma\gamma$, $ZZ^{(*)}$, $WW^{(*)}$, $b\bar{b}$ and $\tau\tau$. only four are independent ZZ and WW related by custodial symmetry
- Loop induced ggF production and $\gamma\gamma$ final state are susceptible to BSM contributions - in case of SUSY light staus and neutralino contribute
- Experimentally we get information on the signal strengths

$$\mu_i \equiv (\sigma \times \text{BR})_i / (\sigma \times \text{BR})_i^{\text{SM}},$$


for each final state

126 GeV Higgs at the LHC


- A combined likelihood in (ggF+ttH) and (VBF+VH) planes was derived using ATLAS, CMS and Tevatron results



- How much invisible Higgs decay is allowed?



- Best fit at zero, for simple SM + invisible case $BR_{inv} < 0.19$
- When all the couplings are allowed to vary $BR_{inv} < 0.38$
- Plenty of room for invisible Higgs decay



A real life application


Do LHC results on the SUSY particles, Higgs signal strengths and constraints on DM from direct and indirect detection experiments rule out light neutralino DM?

Already many studies exist in literature, I'll not list them here

How light is light?

- Relaxing gaugino universality: few collider constraints
 - Z width, LEP bounds, invisible Higgs decays



- Cosmology constraints:



- Region of interest: $m_{\tilde{\chi}_1^0} < m_h/2$ - Z and H exchange not effective, slepton exchange channel has collider constraints from LHC searches
- Light slepton exchange of interest to us here



How light is light?

- pMSSM scan over 11 parameters
 - $M_1, M_2, \mu, \tan\beta, M_A, A_t, M_{1L}, M_{1R}, M_{3L}, M_{3R}, A_\tau$
 - LEP limits, Z width, B-physics, Higgs mass + couplings, heavy Higgs searches @LHC, Xenon100



How light is light?




- DM < 35 GeV associated with light stau + light chargino




- LHC searches put constraints on light electroweak -ino and slepton production

LHC searches

● Direct electroweak -ino production





Applying SModelS



- SMS results used from ATLAS-CONF-2013-049, CMS-PAS-SUS-12-022, ATLAS-CONF-2013-035
- Density of points reduced - LHC does rule out some scenarios
- In general light neutralino still possible

Higgs signal strengths



- Lightest neutralino associated with some invisible Higgs decays
- At 14 TeV with $ZH \rightarrow$ invisible, better sensitivity expected

SMS approach - what's next?

- SMS approach is not perfect yet
- Not all SMS topologies are present
- Need more information from experiments
- Likelihood information is missing which can be used to combine different SMS results, in addition efficiency maps should be built
- Forces are being joined in order to carry out these tasks

Coordinating a simplified models effort

29-30 October 2013
CERN
Europe/Zurich timezone

- Overview
- Timetable
- Registration
 - Registration Form
- List of registrants
- Video Services

Simplified Models have become one of the standard methods to search for physics beyond the Standard Model, reducing full models with dozens of particles and hundreds of parameters down to a handful of particles and parameters, allowing for the commonalities between different models to become apparent.

The challenge with Simplified Models is that any full model decomposes into many different Simplified Models. The spectrum of Simplified Models sometimes requires several different particles with many different parameters. Existing experimental analyses are often sensitive to these alternate Simplified Models, so there is not necessarily any additional work for the experiments to do in the limit setting process, it is only a matter of reinterpreting existing results. While the experimental collaborations can do this, it is often beyond their computing resources and the effort required in reinterpreting results could be spent in performing new analyses.

There are multiple reasons that coordination is desirable. First, experimental analyses often present results on related, but different Simplified Models. Next, there is no standard format to present the results from Simplified Model results. Finally, there are too many Simplified Models for every relevant experimental analysis to present efficiencies for every Simplified Model.

This Workshop aims at stimulating more coordination of efforts between theoretical groups and between experimental efforts, starting with the formation of a Simplified Model Working Group. The goals of this workshop are to:



A. Weiler, J. Wacker, M. Kramer, P. Bechtle, M. Papucci, MadAnalysis, SModelS etc.

Conclusions

- SMS results are a good way to test BSM theories and can have a good constraining power
- SModelS is designed to utilize this power and constrain BSM scenarios
- The formalism is generic and can be applied to any BSM spectra for which SMS results are applicable
- Light neutralino dark matter still compatible with all the direct, indirect detection constraints and LHC searches (LHC searches tested by SModelS)
- Light neutralinos will further be tested by Higgs invisible decays at 14 TeV
- There is still room for improvement for SModelS

Back-up

Indirect detection limits



- We test for FERMI-LAT limits - photons produced from DM annihilation in dwarf spheroidal galaxies in bbar or tautau channel
- Large fraction of LSP < 30 GeV points are several orders of magnitude below the limit

Why not use monojet channel?

- Direct LSP production probed via monojet signature at the LHC
- Limits given on the spin-independent interactions of DM
- Limits applicable for models involving heavy mediators
- Not applicable for MSSM since the mediators are not heavy enough


Scan details

$\tan \beta$	[5, 50]	M_{L_3}	[70, 500]
M_{A^0}	[100, 1000]	M_{R_3}	[70, 500]
M_1	[10, 70]	A_τ	[-1000, 1000]
M_2	[100, 1000]	M_{L_1}	[100, 500]
μ	[100, 1000]	M_{R_1}	[100, 500]


LEP limits	$m_{\tilde{\chi}_1^\pm} > 100 \text{ GeV}$ $m_{\tilde{\tau}_1} > 84 - 88 \text{ GeV}$ (depending on $m_{\tilde{\chi}_1^0}$) $\sigma(e^+e^- \rightarrow \tilde{\chi}_{2,3}^0 \tilde{\chi}_1^0 \rightarrow Z^{(*)}(\rightarrow q\bar{q})\tilde{\chi}_1^0) \lesssim 0.05 \text{ pb}$
invisible Z decay	$\Gamma_{Z \rightarrow \tilde{\chi}_1^0 \tilde{\chi}_1^0} < 3 \text{ MeV}$
μ magnetic moment	$\Delta a_\mu < 4.5 \times 10^{-9}$
flavor constraints	$\text{BR}(b \rightarrow s\gamma) \in [3.03, 4.07] \times 10^{-4}$ $\text{BR}(B_s \rightarrow \mu^+\mu^-) \in [1.5, 4.3] \times 10^{-9}$
Higgs mass	$m_{h^0} \in [122.5, 128.5] \text{ GeV}$
$A^0, H^0 \rightarrow \tau^+\tau^-$	CMS results for $\mathcal{L} = 17 \text{ fb}^{-1}$, m_h^{max} scenario
Higgs couplings	ATLAS, CMS and Tevatron global fit, see text
relic density	$\Omega h^2 < 0.131$ or $\Omega h^2 \in [0.107, 0.131]$
direct detection	XENON100 upper limit
indirect detection	Fermi-LAT bound on gamma rays from dSphs
$pp \rightarrow \tilde{\chi}_2^0 \tilde{\chi}_1^\pm$ $pp \rightarrow \tilde{\ell}^+ \tilde{\ell}^-$	Simplified Models Spectra approach, see text

Tau dominated scenario



- For topologies involving an intermediate particles, three mass slices are given. We interpolate over these slices



interpolate using
k-factors



interpolate and
derive k-factors



interpolate using
k-factors



SMS approach - what's next?

- SMS approach is not perfect yet
- Not all SMS topologies are present
- Need more information from experiments
- Likelihood information is missing which can be used to combine different SMS results, in addition efficiency maps should be built
- Forces are being joined in order to carry out these tasks

The screenshot shows a website for a workshop. The header is a dark blue bar with the title 'Coordinating a simplified models effort' in white. Below the header, the dates '29-30 October 2013' and 'CERN' are listed, along with the location 'Europe/Zurich timezone'. A search bar is visible on the right. A navigation menu on the left includes 'Overview', 'Timetable', 'Registration', 'Registration Form', 'List of registrants', and 'Video Services'. The main content area contains three paragraphs of text discussing the challenges and goals of the workshop.

Coordinating a simplified models effort

29-30 October 2013
CERN
Europe/Zurich timezone

Search

Overview

- Timetable
- Registration
- Registration Form
- List of registrants
- Video Services

Simplified Models have become one of the standard methods to search for physics beyond the Standard Model, reducing full models with dozens of particles and hundreds of parameters down to a handful of particles and parameters, allowing for the commonalities between different models to become apparent.

The challenge with Simplified Models is that any full model decomposes into many different Simplified Models. The spectrum of Simplified Models sometimes requires several different particles with many different parameters. Existing experimental analyses are often sensitive to these alternate Simplified Models, so there is not necessarily any additional work for the experiments to do in the limit setting process, it is only a matter of reinterpreting existing results. While the experimental collaborations can do this, it is often beyond their computing resources and the effort required in reinterpreting results could be spent in performing new analyses.

There are multiple reasons that coordination is desirable. First, experimental analyses often present results on related, but different Simplified Models. Next, there is no standard format to present the results from Simplified Model results. Finally, there are too many Simplified Models for every relevant experimental analysis to present efficiencies for every Simplified Model.

This Workshop aims at stimulating more coordination of efforts between theoretical groups and between experimental efforts, starting with the formation of a Simplified Model Working Group. The goals of this workshop are to: