Structure of the WDW equation

Diffeomorphism invariance and emergence of probabilistic interpretation in QC

Renaud Parentani[1]

[1]LPT, Paris-Sud Orsay

SISSA, September 2011

Based on: PRD56, NPB513, PRD59 (1998-99) with S. Massar and conversations (2005) with T. Jacobson

The context

- There is a **probabilistic interpretation** of the solutions of the Schroedinger eq., but there is **no consensus** on the **interpretation** of the solutions of the WDW eq.

- Several options have been proposed:
  - **"Page-Hawking"** based on the **norm**: $|\Psi(a,\phi)|^2$
  - **"Vilenkin"** based on the **current**: $W = \Psi^* i \overleftrightarrow{\partial_a} \Psi$
  - Third quantization.

  - Square root approach $i\partial_X \Psi = H\Psi$ (Ashtekar (2008))
  - Adding "dust" (Timan (2006))

The context

- There is a **probabilistic interpretation** of the solutions of the Schroedinger eq., but there is **no consensus** on the **interpretation** of the solutions of the WDW eq.

- Several options have been proposed:
  - **"Page-Hawking"** based on the **norm**: $|\Psi(a,\phi)|^2$
  - **"Vilenkin"** based on the **current**: $W = \Psi^* i \overleftrightarrow{\partial_a} \Psi$
  - Third quantization.

  - Square root approach $i\partial_X \Psi = H\Psi$ (Ashtekar (2008))
  - Adding "dust" (Timan (2006))

The context

- There is a **probabilistic interpretation** of the solutions of the Schroedinger eq., but there is **no consensus** on the **interpretation** of the solutions of the WDW eq.

- Several options have been proposed:
  - **"Page-Hawking"** based on the **norm**: $|\Psi(a,\phi)|^2$
  - **"Vilenkin"** based on the **current**: $W = \Psi^* i \overleftrightarrow{\partial_a} \Psi$
  - Third quantization.

  - Square root approach $i\partial_X \Psi = H\Psi$ (Ashtekar (2008))
  - Adding "dust" (Timan (2006))

Plan

- **Compare** (mathem.) the *structure* of the Schr. eq. to that of the WdW eq.
- Focus on **Transition Amplitudes** and use **molecular physics techniques** as a **tool** to perform the comparison.
- Establish that
  - no **exact** proba. interp. could possibly be given: **Diffeo-invar.** + **canon. quantization** $\rightarrow$ no proba. interp.
  - the **Schrod. probabilistic interpretation** is an **emergent** property of the sol. of the WDW eq.

Plan

- **Compare** (mathem.) the *structure* of the Schr. eq. to that of the WdW eq.
- Focus on **Transition Amplitudes** and use **molecular physics techniques** as a **tool** to perform the comparison.
- Establish that
  - no **exact** proba. interp. could possibly be given: **Diffeo-invar.** + **canon. quantization** $\rightarrow$ no proba. interp.
  - the **Schrod. probabilistic interpretation** is an **emergent** property of the sol. of the WDW eq.

Plan

- **Compare** (mathem.) the *structure* of the Schr. eq. to that of the WdW eq.
- Focus on **Transition Amplitudes** and use **molecular physics techniques** as a **tool** to perform the comparison.
- Establish that
  - no **exact** proba. interp. could possibly be given: **Diffeo-invar.** + **canon. quantization** $\rightarrow$ no proba. interp.
  - the **Schrod. probabilistic interpretation** is an **emergent** property of the sol. of the WDW eq.

Why focusing on Transition Amplitudes ?

- The WdW eq. *also* describes **matter** transitions,
  e.g. $e^+ e^-$ annihilation.
  $\rightarrow$ How to **compute** their amplitudes ?

- Historically, **Born's statistical interpretation (1926)**

$$\begin{aligned} |n_0\rangle &\rightarrow \sum_n c_{n,n_0} \, |n\rangle \\ |c_{n,n_0}|^2 &= \text{Proba. to find } |n\rangle \text{ at late time,} \\ &\quad \text{when starting from } |n_0\rangle. \end{aligned} \tag{1}$$

  followed from his understanding of **Transition Amplit**.

- Follow here the same logic:
  study the properties of $\mathcal{C}_n(a)$, **Trans. Amplit. in QC**,
  then consider their interpretation.

Why focusing on Transition Amplitudes ?

- The WdW eq. *also* describes **matter** transitions, *e.g.* $e^+ e^-$ annihilation.
  $\rightarrow$ How to **compute** their amplitudes ?

- Historically, **Born's statistical interpretation (1926)**

$$
\begin{aligned}
|n_0\rangle &\rightarrow \sum_n c_{n,n_0} \, |n\rangle \\
|c_{n,n_0}|^2 &= \text{Proba. to find } |n\rangle \text{ at late time,} \\
&\quad \text{when starting from } |n_0\rangle.
\end{aligned}
\tag{1}
$$

  followed from his understanding of **Transition Amplit**.

- Follow here the same logic:
  study the properties of $\mathcal{C}_n(a)$, **Trans. Amplit. in QC**,
  then consider their interpretation.

Why focusing on Transition Amplitudes ?

- The WdW eq. *also* describes **matter** transitions, *e.g.* $e^+ e^-$ annihilation.
  $\rightarrow$ How to **compute** their amplitudes ?

- Historically, **Born's statistical interpretation (1926)**

$$\begin{aligned} |n_0\rangle &\rightarrow \sum_n c_{n,n_0} \, |n\rangle \\ |c_{n,n_0}|^2 &= \text{Proba. to find } |n\rangle \text{ at late time,} \\ &\quad \text{when starting from } |n_0\rangle. \end{aligned} \tag{1}$$

  followed from his understanding of **Transition Amplit**.

- Follow here the same logic:
  study the properties of $\mathcal{C}_n(a)$, **Trans. Amplit. in QC**,
  then consider their interpretation.

Why focusing on Transition Amplitudes ?

- The WdW eq. *also* describes **matter** transitions, *e.g.* $e^+ e^-$ annihilation.
  $\rightarrow$ How to **compute** their amplitudes ?

- Historically, **Born's statistical interpretation (1926)**

$$\begin{aligned} |n_0\rangle &\rightarrow \sum_n c_{n,n_0} \, |n\rangle \\ |c_{n,n_0}|^2 &= \text{Proba. to find } |n\rangle \text{ at late time,} \\ &\quad \text{when starting from } |n_0\rangle. \end{aligned} \tag{1}$$

  followed from his understanding of **Transition Amplit**.

- Follow here the same logic:
  study the properties of $\mathcal{C}_n(a)$, **Trans. Amplit. in QC**,
  then consider their interpretation.

Why using *molecular physics techniques* ?

- Appropriate for dynamical systems containing **light** and **heavy** (i.e. fast and slow) deg. of freedom.
- **Transition amplitudes** are governed by **frequency ratios** and not by *coupling constants* (called *Non-Adiabatic Transition Amplitudes* **NATA**)
- certain NATA are **exponentially suppressed** w.r.t. others, thereby introducing a **hierarchy of NATA**.
- **use** this **hierarchy** to
  - **organize** solutions of the WDW eq.
  - get **algebraic relations** between Schrod's $c_n(t)$ and WDW's $\mathcal{C}_n(a)$.
- *Long tradition*: Born '26, Heisenberg '35, Gottfried '66 + '98
  All considering the statistical interpret. of QM.

Why using *molecular physics techniques* ?

- Appropriate for dynamical systems containing **light** and **heavy** (i.e. fast and slow) deg. of freedom.
- **Transition amplitudes** are governed by **frequency ratios** and not by *coupling constants* (called *Non-Adiabatic Transition Amplitudes* **NATA**)
- certain NATA are **exponentially suppressed** w.r.t. others, thereby introducing a **hierarchy of NATA**.
- **use** this **hierarchy** to
  - **organize** solutions of the WDW eq.
  - get **algebraic relations** between Schrod's $c_n(t)$ and WDW's $\mathcal{C}_n(a)$.
- *Long tradition*: Born '26, Heisenberg '35, Gottfried '66 + '98
  All considering the statistical interpret. of QM.

Why using *molecular physics techniques* ?

- Appropriate for dynamical systems containing **light** and **heavy** (i.e. fast and slow) deg. of freedom.
- **Transition amplitudes** are governed by **frequency ratios** and not by *coupling constants* (called *Non-Adiabatic Transition Amplitudes* **NATA**)
- certain NATA are **exponentially suppressed** w.r.t. others, thereby introducing a **hierarchy of NATA**.
- **use** this **hierarchy** to
  - **organize** solutions of the WDW eq.
  - get **algebraic relations** between Schrod's $c_n(t)$ and WDW's $\mathcal{C}_n(a)$.
- *Long tradition*: Born '26, Heisenberg '35, Gottfried '66 + '98
  All considering the statistical interpret. of QM.

Why using *molecular physics techniques* ?

- Appropriate for dynamical systems containing **light** and **heavy** (i.e. fast and slow) deg. of freedom.
- **Transition amplitudes** are governed by **frequency ratios** and not by *coupling constants* (called *Non-Adiabatic Transition Amplitudes* **NATA**)
- certain NATA are **exponentially suppressed** w.r.t. others, thereby introducing a **hierarchy of NATA**.
- **use** this **hierarchy** to
  - **organize** solutions of the WDW eq.
  - get **algebraic relations** between Schrod's $c_n(t)$ and WDW's $\mathcal{C}_n(a)$.
- *Long tradition*: Born '26, Heisenberg '35, Gottfried '66 + '98
  All considering the statistical interpret. of QM.

Why using *molecular physics techniques* ?

- Appropriate for dynamical systems containing **light** and **heavy** (i.e. fast and slow) deg. of freedom.
- **Transition amplitudes** are governed by **frequency ratios** and not by *coupling constants* (called *Non-Adiabatic Transition Amplitudes* **NATA**)
- certain NATA are **exponentially suppressed** w.r.t. others, thereby introducing a **hierarchy of NATA**.
- **use** this **hierarchy** to
  - **organize** solutions of the WDW eq.
  - get **algebraic relations** between Schrod's $c_n(t)$ and WDW's $\mathcal{C}_n(a)$.
- *Long tradition*: Born '26, Heisenberg '35, Gottfried '66 + '98
  All considering the statistical interpret. of QM.

Why using *molecular physics techniques* ?

- Appropriate for dynamical systems containing **light** and **heavy** (i.e. fast and slow) deg. of freedom.
- **Transition amplitudes** are governed by **frequency ratios** and not by *coupling constants* (called *Non-Adiabatic Transition Amplitudes* **NATA**)
- certain NATA are **exponentially suppressed** w.r.t. others, thereby introducing a **hierarchy of NATA**.
- **use** this **hierarchy** to
  - **organize** solutions of the WDW eq.
  - get **algebraic relations** between Schrod's $c_n(t)$ and WDW's $\mathcal{C}_n(a)$.
- *Long tradition*: Born '26, Heisenberg '35, Gottfried '66 + '98
  All considering the statistical interpret. of QM.

The Schroed. equation in cosmology.

- *For definiteness*,
  consider some matter fields governed by an hermitian H_M in an expanding **compact** RW sp-time described by $a(t)$.
- The S. eq. is
$$i\partial_t|\psi(t)\rangle = H_M\,|\psi(t)\rangle. \tag{2}$$
- Because of the expansion $da/dt > 0$,
$$H_M = H_M(a(t))$$
  is t-dependent. *This is crucial for what follows.*

Structure of the S. eq. in Cosmology. 1

- To reveal its *structure*, and/or to solve it, introduce the **instantaneous eigenstates** of $H_m(a)$:

$$\begin{aligned} H_M(a)\,|\chi_n(a)\rangle &= E_n(a)\,|\chi_n(a)\rangle \\ \langle\chi_n(a)|\chi_m(a)\rangle &= \delta_{n,m} \end{aligned} \tag{3}$$

- decompose $|\psi(t)\rangle$ in this **basis**:

$$|\psi(t)\rangle = \sum_n c_n(t)\, e^{-i\int^t dt' E_n(t')}\, |\chi_n(a(t))\rangle, \tag{4}$$

- compute the $c_n(t)$ by injecting (4) in $i\partial_t|\psi\rangle = H_M\,|\psi\rangle$.

Structure of the S. eq. in Cosmology. 1

- To reveal its *structure*, and/or to solve it, introduce the **instantaneous eigenstates** of $H_m(a)$:

$$\begin{aligned} H_M(a)\,|\chi_n(a)\rangle &= E_n(a)\,|\chi_n(a)\rangle \\ \langle\chi_n(a)|\chi_m(a)\rangle &= \delta_{n,m} \end{aligned} \tag{3}$$

- decompose $|\psi(t)\rangle$ in this **basis**:

$$|\psi(t)\rangle = \sum_n c_n(t)\; e^{-i\int^t dt' E_n(t')}\; |\chi_n(a(t))\rangle, \tag{4}$$

- compute the $c_n(t)$ by injecting (4) in $i\partial_t|\psi\rangle = H_M\;|\psi\rangle$.

Structure of the S. eq. in Cosmology. 2

This gives a $N \times N$ matricial eq.

$$\partial_t c_n = \sum_m \langle \partial_t \chi_m | \chi_n \rangle \, e^{-i \int^t dt' (E_m - E_n)} c_m. \tag{5}$$

where

$$\langle \chi_m | \partial_t \chi_n \rangle = \frac{\langle \chi_m | \partial_t H_M | \chi_n \rangle}{E_n(a) - E_m(a)} \quad , \quad n \neq m \tag{6}$$

NB. When $da/dt = 0$, one has

$$\partial_t c_n = 0.$$

Hence all NA transitions are induced by da/dt.

Structure of the S. eq. in Cosmology. 2

This gives a $N \times N$ matricial eq.

$$\partial_t c_n = \sum_m \langle \partial_t \chi_m | \chi_n \rangle \, e^{-i \int^t dt' (E_m - E_n)} c_m. \tag{5}$$

where

$$\langle \chi_m | \partial_t \chi_n \rangle = \frac{\langle \chi_m | \partial_t H_M | \chi_n \rangle}{E_n(a) - E_m(a)} \quad , \quad n \neq m \tag{6}$$

NB. When $da/dt = 0$, one has

$$\partial_t c_n \equiv 0.$$

Hence all NA transitions are induced by da/dt.

Structure of the S. eq. in Cosmology. 2 Remarks.

- The amplitudes $d_n(a)$ defined in the **corresponding contracting** universe, $da/dt \to -da/dt$, obey the **same** equation with $i \to -i$.

- The Schrod. eq. **can** be written in terms of a:

$$\partial_a c_n = \sum_{m \neq n} \langle \partial_a \chi_m | \chi_n \rangle \, e^{-i \int^a da' (dt/da')(E_m - E_n)} \, c_m(a). \qquad (7)$$

The cosmic time t **only** appears through $t(a)$.

Structure of the S. eq. in Cosmology. 2 Remarks.

- The amplitudes $d_n(a)$ defined in the **corresponding contracting** universe, $da/dt \to -da/dt$, obey the **same** equation with $i \to -i$.

- The Schrod. eq. **can** be written in terms of a:

$$\partial_a c_n = \sum_{m \neq n} \langle \partial_a \chi_m | \chi_n \rangle \, e^{-i \int^a da' (dt/da')(E_m - E_n)} \, c_m(a). \qquad (7)$$

The cosmic time t **only** appears through $t(a)$.

Structure of the S. eq. in Cosmology. 2 Remarks.

- The amplitudes $d_n(a)$ defined in the **corresponding contracting** universe, $da/dt \to -da/dt$, obey the **same** equation with $i \to -i$.

- The Schrod. eq. **can** be written in terms of a:

$$\partial_a c_n = \sum_{m \neq n} \langle \partial_a \chi_m | \chi_n \rangle \, e^{-i \int^a da' (dt/da')(E_m - E_n)} \, c_m(a). \qquad (7)$$

The cosmic time t **only** appears through $t(a)$.

NATA: Non-Adiabatic Transition Amplitudes.

- Starting with $c_1(t=-\infty)=1$, the amplitude to find the system in the state n is $c_n(+\infty)$.
- To first order in NA, it is

$$c_n(+\infty) \simeq \int_{-\infty}^{+\infty} dt \, \langle \chi_n | \partial_t \chi_1 \rangle \, e^{-i \int_{-\infty}^{t} dt' (E_1(t') - E_n(t'))} \tag{8}$$

- When $c_n \ll 1$, can be evaluated by a saddle point approx.
- The sp time t^* is complex, and hence c_n is expon. damped:

$$c_n(+\infty) \simeq C \, e^{-i \int_{-\infty}^{t^*} dt' (E_1 - E_n)}$$

NB. $C \to 1$ in the adiabatic limit, see refs.

NATA: Non-Adiabatic Transition Amplitudes.

- Starting with $c_1(t=-\infty)=1$, the amplitude to find the system in the state n is $c_n(+\infty)$.
- To first order in NA, it is

$$c_n(+\infty) \simeq \int_{-\infty}^{+\infty} dt \langle \chi_n | \partial_t \chi_1 \rangle \; e^{-i \int_{-\infty}^{t} dt' (E_1(t') - E_n(t'))} \tag{8}$$

- When $c_n \ll 1$, can be evaluated by a saddle point approx.
- The sp time t^* is complex, and hence c_n is expon. damped:

$$c_n(+\infty) \simeq C \; e^{-i \int_{-\infty}^{t^*} dt' (E_1 - E_n)}$$

NB. $C \to 1$ in the adiabatic limit, see refs.

WDW Equation

- We now replace $[-i\partial_t + H_M]\,|\psi(t)\rangle = 0$ by the WDW eq.

$$[H_G + H_M]\,|\Psi(a)\rangle = 0 \tag{9}$$

where

$$H_G = \frac{-G^2\pi_a^2 - a^2 + \Lambda a^4}{2Ga}. \tag{10}$$

$\hat{\pi}_a = -i\partial_a$ is the momentum conjugated to a.

- With (9, 10) we have displaced the *Heisenberg cut* so as to **include** (a, π_a) in the quantum description.

- Eqs. (9, 10) follow from $S = \int d^4x\sqrt{g}R + L_M$ when
  - 3 geometries are compact, and
  - matter distribution is sufficiently homogeneous.

N.B. $H_G + H_M = 0$ **is the Friedmann eq.**

WDW Equation

- We now replace $[-i\partial_t + H_M]\,|\psi(t)\rangle = 0$ by the WDW eq.

$$[H_G + H_M]\,|\Psi(a)\rangle = 0 \tag{9}$$

where

$$H_G = \frac{-G^2\pi_a^2 - a^2 + \Lambda a^4}{2Ga}. \tag{10}$$

$\hat{\pi}_a = -i\partial_a$ is the momentum conjugated to a.

- With (9, 10) we have displaced the *Heisenberg cut* so as to **include** (a, π_a) in the quantum description.

- Eqs. (9, 10) follow from $S = \int d^4x\sqrt{g}R + L_M$ when
  - 3 geometries are compact, and
  - matter distribution is sufficiently homogeneous.

N.B. $H_G + H_M = 0$ **is the Friedmann eq.**

WDW Equation

- We now replace $[-i\partial_t + H_M]\,|\psi(t)\rangle = 0$ by the WDW eq.

$$[H_G + H_M]\,|\Psi(a)\rangle = 0 \tag{9}$$

where

$$H_G = \frac{-G^2\pi_a^2 - a^2 + \Lambda a^4}{2Ga}. \tag{10}$$

$\hat{\pi}_a = -i\partial_a$ is the momentum conjugated to a.

- With (9, 10) we have displaced the *Heisenberg cut* so as to **include** (a, π_a) in the quantum description.

- Eqs. (9, 10) follow from $S = \int d^4x\sqrt{g}R + L_M$ when
  - 3 geometries are compact, and
  - matter distribution is sufficiently homogeneous.

N.B. $H_G + H_M = 0$ **is the Friedmann eq.**

Conserved Wronskian

- The **unique** conserved quantity associated with the Schrod. eq. is the **norm**

$$N = \langle \psi(t) | \psi(t) \rangle \equiv Cst. = \sum_n |c_n(t)|^2. \tag{11}$$

- **Instead** the **unique** one associated with the WDW eq. is the **Wronskian**

$$W = \langle \Psi(a) | \, i \overleftrightarrow{\partial_a} | \Psi(a) \rangle \equiv Cst. \tag{12}$$

- Question: **Can** W **be written as** $W = \sum_n |\mathcal{C}_n(a)|^2$ **?**

Conserved Wronskian

- The **unique** conserved quantity associated with the Schrod. eq. is the **norm**

$$N = \langle \psi(t) | \psi(t) \rangle \equiv Cst. = \sum_n |c_n(t)|^2. \tag{11}$$

- **Instead** the **unique** one associated with the WDW eq. is the **Wronskian**

$$W = \langle \Psi(a) | \, i \overleftrightarrow{\partial_a} | \Psi(a) \rangle \equiv Cst. \tag{12}$$

- Question: **Can W be written as $W = \sum_n |\mathcal{C}_n(a)|^2$?**

Conserved Wronskian

- The **unique** conserved quantity associated with the Schrod. eq. is the **norm**

$$N = \langle \psi(t) | \psi(t) \rangle \equiv Cst. = \sum_n |c_n(t)|^2. \tag{11}$$

- **Instead** the **unique** one associated with the WDW eq. is the **Wronskian**

$$W = \langle \Psi(a) | \, i \overleftrightarrow{\partial_a} | \Psi(a) \rangle \equiv Cst. \tag{12}$$

- Question: **Can** W **be written as** $W = \sum_n |\mathcal{C}_n(a)|^2$ **?**

First attempt

- Decompose

$$|\Psi(a)\rangle = \sum_n \mathcal{C}_n(a)\, \Psi_n(a)\, |\chi_n(a)\rangle, \tag{13}$$

where $|\chi_n(a)\rangle$ is the **same** as for the Schrod. eq.
and $\Psi_n(a)$ is the WKB solution of

$$[H_G + E_n(a)]\, \Psi_n(a) = 0 \tag{14}$$

- The unit positive Wronskian WKB solution is

$$\Psi_n(a) = \frac{e^{-i \int^a da\; p_n(a)}}{\sqrt{2p_n(a)}} \tag{15}$$

where the momentum $p_n(a) > 0$ solves $H_G + E_n(a) = 0$.
These solutions correspond to expanding universes.
Their complex conjugated describe **contracting** universes.

First attempt

- Decompose

$$|\Psi(a)\rangle = \sum_n \mathcal{C}_n(a)\, \Psi_n(a)\, |\chi_n(a)\rangle, \tag{13}$$

where $|\chi_n(a)\rangle$ is the **same** as for the Schrod. eq.
and $\Psi_n(a)$ is the WKB solution of

$$[H_G + E_n(a)]\, \Psi_n(a) = 0 \tag{14}$$

- The unit positive Wronskian WKB solution is

$$\Psi_n(a) = \frac{e^{-i \int^a da\, p_n(a)}}{\sqrt{2p_n(a)}} \tag{15}$$

where the momentum $p_n(a) > 0$ solves $H_G + E_n(a) = 0$.

These solutions correspond to expanding universes.

Their complex conjugated describe **contracting** universes.

First attempt

- inserting (13) in the WDW eq. gives a **second** order matricial equation which mixes
  - corrections to WKB corrections with
  - $n \to n'$ matter transitions.
- Moreover the conserved Wronskian W reads

$$W = \sum_n |\mathcal{C}_n(a)|^2 + \sum_n (\mathcal{C}_n^* i \overleftrightarrow{\partial_a} \mathcal{C}_n) \, |\Psi_n(a)|^2. \tag{16}$$

The first term is OK, the second unwanted.

- Question: How to sort this out ?

First attempt

- inserting (13) in the WDW eq. gives a **second** order matricial equation which mixes
  - corrections to WKB corrections with
  - $n \to n'$ matter transitions.
- Moreover the conserved Wronskian W reads

$$W = \sum_n |\mathcal{C}_n(a)|^2 + \sum_n (\mathcal{C}_n^* i \overleftrightarrow{\partial_a} \mathcal{C}_n) \, |\Psi_n(a)|^2. \tag{16}$$

The first term is OK, the second unwanted.

- Question: How to sort this out ?

First attempt

- inserting (13) in the WDW eq. gives a **second** order matricial equation which mixes
  - corrections to WKB corrections with
  - $n \to n'$ matter transitions.
- Moreover the conserved Wronskian W reads

$$W = \sum_n |\mathcal{C}_n(a)|^2 + \sum_n (\mathcal{C}_n^* i \overleftrightarrow{\partial_a} \mathcal{C}_n) \, |\Psi_n(a)|^2. \tag{16}$$

  The first term is OK, the second unwanted.

- Question: How to sort this out ?

WDW equation: The correct decomposition

- The "correct" decomposition

$$|\Psi(a)\rangle = \sum_n \left[\mathcal{C}_n(a)\ \Psi_n(a) + \mathcal{D}_n(a)\ \Psi_n^*(a)\right]\ |\chi_n(a)\rangle, \qquad (17)$$

  introduces $2N$ arbitrary fonctions $\mathcal{C}_n(a), \mathcal{D}_n(a)$.

- Eq. (17) gives an **under-constrained** system.
- **Exploit** this and impose

$$\langle\chi_n|i\overrightarrow{\partial_a}|\Psi\rangle = p_n\left[\mathcal{C}_n\ \Psi_n - \mathcal{D}_n\ \Psi_n^*\right]. \qquad (18)$$

  Now the $\mathcal{C}_n$ ($\mathcal{D}_n$) **instantanenously** weigh expanding (contracting) solutions.

- Insert (17) in the WdW eq. *and* use (18) to get

WDW equation: The correct decomposition

- The "correct" decomposition

$$|\Psi(a)\rangle = \sum_n \left[\mathcal{C}_n(a)\ \Psi_n(a) + \mathcal{D}_n(a)\ \Psi_n^*(a)\right]\ |\chi_n(a)\rangle, \qquad (17)$$

  introduces $2N$ arbitrary fonctions $\mathcal{C}_n(a), \mathcal{D}_n(a)$.

- Eq. (17) gives an **under-constrained** system.

- **Exploit** this and impose

$$\langle\chi_n|i\overrightarrow{\partial_a}|\Psi\rangle = p_n\left[\mathcal{C}_n\ \Psi_n - \mathcal{D}_n\ \Psi_n^*\right]. \qquad (18)$$

  Now the $\mathcal{C}_n$ ($\mathcal{D}_n$) **instantanenously** weigh expanding (contracting) solutions.

- Insert (17) in the WdW eq. *and* use (18) to get

WDW equation: The correct decomposition

- The "correct" decomposition

$$|\Psi(a)\rangle = \sum_n \left[\mathcal{C}_n(a)\ \Psi_n(a) + \mathcal{D}_n(a)\ \Psi_n^*(a)\right]\ |\chi_n(a)\rangle, \qquad (17)$$

  introduces $2N$ arbitrary fonctions $\mathcal{C}_n(a), \mathcal{D}_n(a)$.

- Eq. (17) gives an **under-constrained** system.
- **Exploit** this and impose

$$\langle\chi_n|i\overrightarrow{\partial_a}|\Psi\rangle = p_n\left[\mathcal{C}_n\ \Psi_n - \mathcal{D}_n\ \Psi_n^*\right]. \qquad (18)$$

  Now the $\mathcal{C}_n$ ($\mathcal{D}_n$) **instantanenously** weigh expanding (contracting) solutions.

- Insert (17) in the WdW eq. *and* use (18) to get

WDW equation: The correct decomposition

- The "correct" decomposition

$$|\Psi(a)\rangle = \sum_n \left[\mathcal{C}_n(a)\ \Psi_n(a) + \mathcal{D}_n(a)\ \Psi_n^*(a)\right]\ |\chi_n(a)\rangle, \qquad (17)$$

  introduces $2N$ arbitrary fonctions $\mathcal{C}_n(a), \mathcal{D}_n(a)$.

- Eq. (17) gives an **under-constrained** system.
- **Exploit** this and impose

$$\langle\chi_n|i\overrightarrow{\partial_a}|\Psi\rangle = p_n\left[\mathcal{C}_n\ \Psi_n - \mathcal{D}_n\ \Psi_n^*\right]. \qquad (18)$$

  Now the $\mathcal{C}_n$ ($\mathcal{D}_n$) **instantanenously** weigh expanding (contracting) solutions.

- Insert (17) in the WdW eq. *and* use (18) to get

Structure of the WDW equation, 1

$$\partial_a \mathcal{C}_n = \sum_{m\neq n} \tilde{M}_{nm}\, e^{-i\int^a (p_n - p_m) da}\, \mathcal{C}_m + \sum_m \tilde{N}_{nm}\, e^{-i\int^a (p_n + p_m) da}\, \mathcal{D}_m. \tag{19}$$

+ the same eq. with $\mathcal{C}_n \to \mathcal{D}_n$ and $i \to -i$
where

$$\tilde{M}_{nm} = \langle \partial_a \psi_m | \psi_n \rangle \frac{p_n + p_m}{2\sqrt{p_n p_m}} \tag{20}$$

$$\tilde{N}_{nm} = \langle \partial_a \psi_m | \psi_n \rangle \frac{p_n - p_m}{2\sqrt{p_n p_m}} + \delta_{nm} \frac{\partial_a p_n}{2 p_n} \tag{21}$$

- Eqs. (19,20,21) are **exact**
- **Eq.(19) is the WDW eq.**

Structure of the WDW equation, 1

$$\partial_a \mathcal{C}_n = \sum_{m\neq n} \tilde{M}_{nm}\, e^{-i\int^a (p_n - p_m) da}\, \mathcal{C}_m + \sum_m \tilde{N}_{nm}\, e^{-i\int^a (p_n + p_m) da}\, \mathcal{D}_m. \quad (19)$$

+ the same eq. with $\mathcal{C}_n \to \mathcal{D}_n$ and $i \to -i$
where

$$\tilde{M}_{nm} = \langle \partial_a \psi_m | \psi_n \rangle \frac{p_n + p_m}{2\sqrt{p_n p_m}} \quad (20)$$

$$\tilde{N}_{nm} = \langle \partial_a \psi_m | \psi_n \rangle \frac{p_n - p_m}{2\sqrt{p_n p_m}} + \delta_{nm} \frac{\partial_a p_n}{2 p_n} \quad (21)$$

- Eqs. (19,20,21) are **exact**
- Eq.(19) is the WDW eq.

Structure of the WDW equation, 1

$$\partial_a \mathcal{C}_n = \sum_{m\neq n} \tilde{M}_{nm}\, e^{-i\int^a (p_n - p_m) da}\, \mathcal{C}_m + \sum_m \tilde{N}_{nm}\, e^{-i\int^a (p_n + p_m) da}\, \mathcal{D}_m. \tag{19}$$

+ the same eq. with $\mathcal{C}_n \to \mathcal{D}_n$ and $i \to -i$
where

$$\tilde{M}_{nm} = \langle \partial_a \psi_m | \psi_n \rangle \frac{p_n + p_m}{2\sqrt{p_n p_m}} \tag{20}$$

$$\tilde{N}_{nm} = \langle \partial_a \psi_m | \psi_n \rangle \frac{p_n - p_m}{2\sqrt{p_n p_m}} + \delta_{nm} \frac{\partial_a p_n}{2 p_n} \tag{21}$$

- Eqs. (19,20,21) are **exact**
- **Eq.(19) is the WDW eq.**

Structure of the WDW equation, 2

- Moreover, as an **identity**, one has

$$\begin{aligned} W &= \langle \Psi(a) | i \overleftrightarrow{\partial_a} | \Psi(a) \rangle \\ &= \sum_n |\mathcal{C}_n(a)|^2 - |\mathcal{D}_n(a)|^2 \equiv Const. \end{aligned} \tag{22}$$

  for **any** Hermitian matter hamiltonian H_M,
  and **any** state.

- Let's describe the **consequences** of Eq. 19 and 22

Structure of the WDW equation, 2

- Moreover, as an **identity**, one has

$$\begin{aligned} W &= \langle \Psi(a) | i \overleftrightarrow{\partial_a} | \Psi(a) \rangle \\ &= \sum_n |\mathcal{C}_n(a)|^2 - |\mathcal{D}_n(a)|^2 \equiv Const. \end{aligned} \tag{22}$$

  for **any** Hermitian matter hamiltonian H_M,
  and **any** state.

- Let's describe the **consequences** of Eq. 19 and 22

Theorem 1

- When, for a **given** matter Hamiltonian H_M,
  the Schrod. eq. gives a $N \times N$ first order matricial eq.

  the WDW eq. gives a $2N \times 2N$ first order matricial eq.

- NB. A similar **doubling** is also found in
  non-relativistic molecular/atomic physics.

  It is a general conseq. of QM, when moving the H-cut.

Theorem 1

- When, for a **given** matter Hamiltonian H_M,
  the Schrod. eq. gives a $N \times N$ first order matricial eq.

  the WDW eq. gives a $2N \times 2N$ first order matricial eq.

- NB. A similar **doubling** is also found in
  non-relativistic molecular/atomic physics.

  It is a general conseq. of QM, when moving the H-cut.

Theorem 2

- **Neglecting only** the $\mathcal{C}_n \to \mathcal{D}_n$ transitions, one gets

$$\partial_a \mathcal{C}_n = \sum_{m \neq n} \tilde{M}_{nm}\, e^{-i \int^a (p_n - p_m) da}\, \mathcal{C}_m \tag{23}$$

and

$$\sum_n |\mathcal{C}_n(a)|^2 \equiv Cst.$$

+ the same eq. with $\mathcal{C}_n \to \mathcal{D}_n$ and $i \to -i$.

- WDW eq. thus gives **two** separate (and **unitary**) eqs: one for the expanding sector, one for the contracting.

- Eq. (23) is **not** the Schrod. equation. There is **no** background sp-time, **no** cosmic time (yet)

Theorem 2

- **Neglecting only** the $\mathcal{C}_n \to \mathcal{D}_n$ transitions, one gets

$$\partial_a \mathcal{C}_n = \sum_{m \neq n} \tilde{M}_{nm}\, e^{-i \int^a (p_n - p_m) da}\, \mathcal{C}_m \tag{23}$$

  and

$$\sum_n |\mathcal{C}_n(a)|^2 \equiv Cst.$$

  + the same eq. with $\mathcal{C}_n \to \mathcal{D}_n$ and $i \to -i$.

- WDW eq. thus gives **two** separate (and **unitary**) eqs: one for the expanding sector, one for the contracting.

- Eq. (23) is **not** the Schrod. equation.
  There is **no** background sp-time, **no** cosmic time (yet)

Theorem 2

- **Neglecting only** the $\mathcal{C}_n \to \mathcal{D}_n$ transitions, one gets

$$\partial_a \mathcal{C}_n = \sum_{m \neq n} \tilde{M}_{nm}\, e^{-i \int^a (p_n - p_m) da}\, \mathcal{C}_m \tag{23}$$

  and

$$\sum_n |\mathcal{C}_n(a)|^2 \equiv Cst.$$

  + the same eq. with $\mathcal{C}_n \to \mathcal{D}_n$ and $i \to -i$.

- WDW eq. thus gives **two** separate (and **unitary**) eqs: one for the expanding sector, one for the contracting.

- Eq. (23) is **not** the Schrod. equation.
  There is **no** background sp-time, **no** cosmic time (yet)

Theorem 3. The recovery of cosmic time

- **Neglecting** the $\mathcal{C}_n \to \mathcal{D}_n$ transitions, **and** to **first order** in $E_n - E_{\bar{n}}$ around the mean matter state $\bar{n}$,

$$\mathcal{C}_n(a) \equiv c_n(\bar{t}(a)), \tag{24}$$

where

$$\bar{t}(a) = \int^a da' \partial_E p(a')|_{E=E_{\bar{n}}}, \tag{25}$$

is the HJ time to reach a when matter is in the $\bar{n}$ state.

- The **identity** (24) is Heisenberg-'35 result. It follows from
  - $\tilde{M}_{nm} \equiv M^S_{mn}$
  - $e^{-i\int^a da'(p_n - p_m)} \equiv e^{-i\int^t dt'(E_n - E_m)}$

Theorem 3. The recovery of cosmic time

- **Neglecting** the $\mathcal{C}_n \to \mathcal{D}_n$ transitions, **and** to **first order** in $E_n - E_{\bar{n}}$ around the mean matter state $\bar{n}$,

$$\mathcal{C}_n(a) \equiv c_n(\bar{t}(a)), \tag{24}$$

where

$$\bar{t}(a) = \int^a da' \partial_E p(a')|_{E=E_{\bar{n}}}, \tag{25}$$

is the HJ time to reach a when matter is in the $\bar{n}$ state.

- The **identity** (24) is Heisenberg-'35 result. It follows from
  - $\tilde{M}_{nm} \equiv M^S_{mn}$
  - $e^{-i\int^a da'(p_n - p_m)} \equiv e^{-i\int^t dt'(E_n - E_m)}$

A remark on NATAmplitudes

- For the Schrod. eq., c_n is given by the **overlap**

$$c_n(t) = \langle \chi_n(t) | e^{+i \int^t dt' E_n} | \psi(t) \rangle . \tag{26}$$

Instead, for the WDW eq., $\mathcal{C}_n$ is given by

$$\mathcal{C}_n(a) = \langle \chi_n(a) | \Psi_n(a)^* \, i \overleftrightarrow{\partial_a} | \Psi(a) \rangle \tag{27}$$

This is the *Vilenkin* **current**,
and NOT the *Page-Hawking* overlap $\langle \chi_n(a) | \Psi(a) \rangle$.

A remark on NATAmplitudes

- For the Schrod. eq., c_n is given by the **overlap**

$$c_n(t) = \langle \chi_n(t) | e^{+i \int^t dt' E_n} | \psi(t) \rangle . \tag{26}$$

Instead, for the WDW eq., $\mathcal{C}_n$ is given by

$$\mathcal{C}_n(a) = \langle \chi_n(a) | \Psi_n(a)^* \, i \overleftrightarrow{\partial_a} | \Psi(a) \rangle \tag{27}$$

This is the *Vilenkin* **current**,
and NOT the *Page-Hawking* overlap $\langle \chi_n(a) | \Psi(a) \rangle$.

Theorem 4. The $\mathcal{C}_n \to \mathcal{D}_m$ NATransitions.

- Far from turning points (big bounce), the transitions $\mathcal{C}_n \to \mathcal{D}_m$ are **exponentially suppressed** w.r.t. to the $\mathcal{C}_n \to \mathcal{C}_m$.
- Because, typically

$$\begin{aligned} \mathcal{C}_n \to \mathcal{C}_m &\sim e^{-(p_n - p_m)\, Im \Delta a_C^{sp}} \\ \mathcal{C}_n \to \mathcal{D}_m &\sim e^{-(p_n + p_m)\, Im \Delta a_D^{sp}} \end{aligned} \tag{28}$$

  $p_n + p_m$ scales with the **total matter energy**.
- The Lesson:
  The WDW eq. predicts a hierarchy of NAT, and governs it.

Theorem 4. The $\mathcal{C}_n \to \mathcal{D}_m$ NATransitions.

- Far from turning points (big bounce), the transitions $\mathcal{C}_n \to \mathcal{D}_m$ are **exponentially suppressed** w.r.t. to the $\mathcal{C}_n \to \mathcal{C}_m$.
- Because, typically

$$\begin{aligned} \mathcal{C}_n \to \mathcal{C}_m &\sim e^{-(p_n - p_m)\, Im \Delta a_C^{sp}} \\ \mathcal{C}_n \to \mathcal{D}_m &\sim e^{-(p_n + p_m)\, Im \Delta a_D^{sp}} \end{aligned} \tag{28}$$

  $p_n + p_m$ scales with the **total matter energy**.

- The Lesson:
  The WDW eq. predicts a hierarchy of NAT, and governs it.

Theorem 4. The $\mathcal{C}_n \to \mathcal{D}_m$ NATransitions.

- Far from turning points (big bounce), the transitions $\mathcal{C}_n \to \mathcal{D}_m$ are **exponentially suppressed** w.r.t. to the $\mathcal{C}_n \to \mathcal{C}_m$.
- Because, typically

$$\begin{aligned} \mathcal{C}_n \to \mathcal{C}_m &\sim e^{-(p_n - p_m)\, Im \Delta a_C^{sp}} \\ \mathcal{C}_n \to \mathcal{D}_m &\sim e^{-(p_n + p_m)\, Im \Delta a_D^{sp}} \end{aligned} \tag{28}$$

  $p_n + p_m$ scales with the **total matter energy**.
- The Lesson:
  The WDW eq. predicts a hierarchy of NAT, and governs it.

The $\mathcal{C}_n \to \mathcal{D}_m$ NAT in molecular phys.

- Consider a molecule of mass M and CM R

$$H_{\text{atom}} = \frac{P^2}{2M} + V(R) + H_e(R),$$

where $H_e(R)$ governs the electronic deg. of freedom.

- There are **two** levels of quantization:
- The LZ way: treat R classically as a given function of time $R(t)$. The residual elect. d.o.f. then obey the t-dep. Schrod. eq.

$$i\partial_t|\psi_e\rangle = \hat{H}_e(R(t))|\psi_e\rangle$$

- The WdW way: treat **both** R and el. d.o.f on the **same footing**. Then, at fixed total energy E, one has

$$\left[-\frac{\partial_R^2}{2M} + V(R) + H_{el}(R)\right]|\Psi_E(R)\rangle = E\,|\Psi_E(R)\rangle$$

The $\mathcal{C}_n \to \mathcal{D}_m$ NAT in molecular phys.

- Consider a molecule of mass M and CM R

$$H_{\text{atom}} = \frac{P^2}{2M} + V(R) + H_e(R),$$

where $H_e(R)$ governs the electronic deg. of freedom.

- There are **two** levels of quantization:
- The LZ way: treat R classically as a given function of time $R(t)$. The residual elect. d.o.f. then obey the t-dep. Schrod. eq.

$$i\partial_t|\psi_e\rangle = \hat{H}_e(R(t))|\psi_e\rangle$$

- The WdW way: treat **both** R and el. d.o.f on the **same footing**. Then, at fixed total energy E, one has

$$\left[-\frac{\partial_R^2}{2M} + V(R) + H_{el}(R)\right]|\Psi_E(R)\rangle = E\ |\Psi_E(R)\rangle$$

The $\mathcal{C}_n \to \mathcal{D}_m$ NAT in atomic phys.

Decomposing $|\Psi_E(R)\rangle = \sum_n (\mathcal{C}_n \Psi_n + \mathcal{D}_n \Psi_n^*)|\chi_n(R)\rangle$

- **neglecting** $\mathcal{C}_n \to \mathcal{D}_m$ and to first order in $E_n^e - E_m^e$, one gets

$$\mathcal{C}_n(R) \equiv c_n(\bar{t}(R)),$$

  where c_n are the Landau-Z tr. amplitudes.
  Hence the $|\mathcal{C}_n|^2$ give **transition probablities**.

- The $\mathcal{C}_n(R)$ are given by $\mathcal{C}_n \equiv \langle \chi_n | \Psi_n^* \, i \overleftrightarrow{\partial_R} | \Psi_E \rangle$.,
  i.e. by the "Vilenkin" current.

- Taking into account the NAT $\mathcal{C}_n \to \mathcal{D}_m$, one gets, as an identity,
  $\sum_n |\mathcal{C}_n(R)|^2 - |\mathcal{D}_n(R)|^2 \equiv Const.$
  This is a re-expression of **unitarity**.

- The H-Page norm $|\langle \chi_n | \Psi_E \rangle|^2 = |\mathcal{C}_n \Psi_n(R) + \mathcal{D}_n \Psi_n^*(R)|^2$ is
  - highly interferent, and
  - answers **another** question.

Conclusions. 1

- The WDW eq. determines the NATA.
  There is NO ambiguity in **computing** these amplitudes.

- It predicts that there is a **hierarchy** of NA-regimes.

Conclusions. 2. The S. regime, the mildest one.

- The first regime, the mildest one, is obtained by
  - neglecting $\mathcal{C}_n \to \mathcal{D}_m$ NAT,
  - the first order in $E_n - E_{\bar{n}}$, but **nothing else**.

- In this regime

$$\mathcal{C}_n(a) \equiv c_n(\bar{t}(a)).$$

Therefore $|\mathcal{C}_n(a)|^2$ is the **proba.** to find the state n in an expand. universe at a (in fact around a).

Conclusions. 2. The S. regime, the mildest one, 2.

- Unitarity,

$$\sum_n |\mathcal{C}_n(a)|^2 \equiv Cst.$$

relies on the conserved current W.

- NO problem of interpretation, NO problem of time.

Conclusions. 2. The S. regime, the mildest one, 2.

- Unitarity,

$$\sum_n |\mathcal{C}_n(a)|^2 \equiv Cst.$$

  relies on the conserved current W.

- NO problem of interpretation, NO problem of time.

Conclusions. 3. The second intermediate regime.

- Neglecting **only** $\mathcal{C}_n \to \mathcal{D}_m$, the WDW gives **two separate 1st order eqs. in $i\partial_a$.**

- **Unitarity**, i.e. $\sum_n |\mathcal{C}_n(a)|^2 \equiv Cst.$ is obtained, as in the Schrod. regime, from the conservation of W, even though there is **no cosmic time** because **no** backd sp-time **common** to all matter states.

- Lesson:
  The **recovery of cosmic time** requires more condition than the **recovery of unitarity**.

Conclusions. 3. The second intermediate regime.

- Neglecting **only** $\mathcal{C}_n \to \mathcal{D}_m$, the WDW gives **two separate 1st order eqs. in $i\partial_a$.**

- **Unitarity**, i.e. $\sum_n |\mathcal{C}_n(a)|^2 \equiv Cst.$ is obtained, as in the Schrod. regime, from the conservation of W, even though there is **no cosmic time** because **no** backd sp-time **common** to all matter states.

- Lesson:
  The **recovery of cosmic time** requires more condition than the **recovery of unitarity**.

Conclusions. 3. The second intermediate regime.

- Neglecting **only** $\mathcal{C}_n \to \mathcal{D}_m$, the WDW gives **two separate 1st order eqs. in $i\partial_a$.**

- **Unitarity**, i.e. $\sum_n |\mathcal{C}_n(a)|^2 \equiv Cst.$ is obtained, as in the Schrod. regime, from the conservation of W, even though there is **no cosmic time** because **no** backd sp-time **common** to all matter states.

- Lesson:
  The **recovery of cosmic time** requires more condition than the **recovery of unitarity**.

Conclusions. 4. The least Adiabatic regime.

- Taking into account $\mathcal{C}_n \to \mathcal{D}_m$ NAT, one faces the **identity**

$$\sum_n |\mathcal{C}_n(a)|^2 - |\mathcal{D}_n(a)|^2 \equiv Cst.$$

- How to interprete this equation ?
- *Proposal:* as we just did it for cosmic time:
  the recovery of unitarity requires some conditions.
- Hence we conclude/conjecture:

  The **statistical interpretation** of the NATA $\mathcal{C}_n(a)$
  is **not** as a **fundamental property**,
  but should be conceived as an **emergent property** of QC.

  (Jacobson 2005, Massar & RP 1999, Vilenkin 1989)

Conclusions. 4. The least Adiabatic regime.

- Taking into account $\mathcal{C}_n \to \mathcal{D}_m$ NAT, one faces the **identity**

$$\sum_n |\mathcal{C}_n(a)|^2 - |\mathcal{D}_n(a)|^2 \equiv Cst.$$

- How to interprete this equation ?
- *Proposal:* as we just did it for cosmic time:
  the recovery of unitarity requires some conditions.
- Hence we conclude/conjecture:

  The **statistical interpretation** of the NATA $\mathcal{C}_n(a)$
  is **not** as a **fundamental property**,
  but should be conceived as an **emergent property** of QC.

  (Jacobson 2005, Massar & RP 1999, Vilenkin 1989)

Conclusions. 4. The least Adiabatic regime.

- Taking into account $\mathcal{C}_n \to \mathcal{D}_m$ NAT, one faces the **identity**

$$\sum_n |\mathcal{C}_n(a)|^2 - |\mathcal{D}_n(a)|^2 \equiv Cst.$$

- How to interprete this equation ?
- *Proposal:* as we just did it for cosmic time:
  the recovery of unitarity requires some conditions.
- Hence we conclude/conjecture:

  The **statistical interpretation** of the NATA $\mathcal{C}_n(a)$
  is **not** as a **fundamental property**,
  but should be conceived as an **emergent property** of QC.

  (Jacobson 2005, Massar & RP 1999, Vilenkin 1989)

Conclusions. 4. The least Adiabatic regime.

- Taking into account $\mathcal{C}_n \to \mathcal{D}_m$ NAT, one faces the **identity**

$$\sum_n |\mathcal{C}_n(a)|^2 - |\mathcal{D}_n(a)|^2 \equiv Cst.$$

- How to interprete this equation ?
- *Proposal:* as we just did it for cosmic time:
  the recovery of unitarity requires some conditions.
- Hence we conclude/conjecture:

  The **statistical interpretation** of the NATA $\mathcal{C}_n(a)$
  is **not** as a **fundamental property**,
  but should be conceived as an **emergent property** of QC.

  (Jacobson 2005, Massar & RP 1999, Vilenkin 1989)