


BZ integrals

$$\frac{1}{N_k} \sum_k f(k) = \langle f \rangle_{BZ}$$

$$f(k) = \sum_R f_R e^{+ikR} \Rightarrow \langle f \rangle_{BZ} = f_0$$

a regular grid with $\Delta k = \frac{2\pi}{a} \frac{1}{N}$ integrate exactly up to $R_{MAX} = \frac{Na}{2}$ (in a cubic system)



If some symmetry is present it can be exploited in order to compute only the k -point in the irreducible wedge of the BZ

$$S(R + \tau_s) + f = R' + \tau_s'$$

Monkhorst and Pack
Phys. Rev. B 13 S188 (76)

$$\psi_{Sk, \nu}^*(r) = \psi_{k, \nu}^*(S^{-1}r - f)$$

$\{\vec{R}_i, w_i\}$ with $\sum_i w_i = 1$ and $\vec{R}_i \in I\kappa$

$$\tilde{\rho}(r) = \sum_i w_i |\psi_{k_i, \nu}(r)|^2 \text{ unsymmetrized density}$$

$$\rho(r) = \frac{1}{N_{sym}} \sum_S \tilde{\rho}(S^{-1}r - f)$$

The needed R_{MAX} is of the order of the spatial extent of the Wannier functions of the solid \approx a few bond lengths

For large systems the BZ is small and the bands flat $\Rightarrow \langle f \rangle_{BZ} \approx f(\Gamma)$

SC

$$\{\vec{R}\}$$

$$R^2$$

#

$$\sum_R e^{+iR\kappa} = G_{\Sigma}(\kappa)$$

$$\sum_0 (0,0,0) \quad 0 \quad 1 \quad 1$$

$$\sum_1 (0,0,\pm 1) \quad 1 \quad 6 \quad [\cos(2\pi k_x) + \cos(2\pi k_y) + \cos(2\pi k_z)] \times 2$$

$$\sum_2 (0, \pm 1, \pm 1) \quad 2 \quad 12 \quad [\cos(2\pi k_x) \cdot \cos(2\pi k_y) + \cos(2\pi k_x) \cdot \cos(2\pi k_z) + \cos(2\pi k_y) \cdot \cos(2\pi k_z)]$$

$$\sum_3 (\pm 1, \pm 1, \pm 1) \quad 3 \quad 8 \quad [\cos(2\pi k_x) \cos(2\pi k_y) \cos(2\pi k_z)]$$

⋮
⋮
⋮

$$\begin{aligned} \frac{\sum}{(2\pi)^3} \int d\kappa^3 f(\kappa) &= \frac{\sum}{(2\pi)^3} \int d\kappa^3 \left[\sum_R A(R) e^{+iR\kappa} \right] \\ &= \frac{\sum}{(2\pi)^3} \int d\kappa^3 \left[\sum_{\Sigma} A(\Sigma) \left(\sum_{R \in \Sigma} e^{+iR\kappa} \right) \right] \\ &= \sum_{\Sigma} A(\Sigma) \left(\frac{\sum}{(2\pi)^3} \int d\kappa^3 \sum_{R \in \Sigma} e^{+iR\kappa} \right) = A(\Sigma) \end{aligned}$$

Now $G_{\Sigma_0}(\kappa) = 1 \quad \forall \kappa$

for $\vec{\kappa}^* = \left(\frac{1}{4}, \frac{1}{4}, \frac{1}{4}\right) \therefore G_{\Sigma_1}(\kappa^*) = 0, G_{\Sigma_2}(\kappa^*) = 0, G_{\Sigma_3}(\kappa^*) = 0$

but in general $G_{\Sigma_i}(\kappa^*) \neq 0$ for $i > 3$

fcc

$$\{\vec{R}\} \quad R^2 \quad \# \quad \sum_R e^{+i\vec{R}\cdot\vec{k}} = G_{\Sigma}(\vec{k})$$

$$\sum_0 (0,0,0) \quad 0 \quad 1 \quad 1$$

$$\sum_1 (\pm\frac{1}{2}, \pm\frac{1}{2}, 0) \quad \frac{1}{2} \quad 12 \quad [\cos(k_x\pi)\cos(k_y\pi) + \\ \cos(k_x\pi)\cos(k_z\pi) + \\ \cos(k_y\pi)\cos(k_z\pi)] \cdot 4$$

$$\sum_2 (\pm 1, 0, 0) \quad 1 \quad 6 \quad [\cos(2\pi k_x) + \\ \cos(2\pi k_y) + \\ \cos(2\pi k_z)] \cdot 2$$

$$\sum_3 (\pm 1, \pm \frac{1}{2}, \pm \frac{1}{2}) \quad \frac{3}{2} \quad 24 \quad [\cos(2\pi k_x) \cdot \cos(\pi k_y) \cos(\pi k_z) + \\ \cos(\pi k_x) \cdot \cos(2\pi k_y) \cdot \cos(\pi k_z) + \\ \cos(\pi k_x) \cdot \cos(\pi k_y) \cdot \cos(2\pi k_z)] \cdot 8$$

⋮

NOW

$$G_{\Sigma_0}(\vec{k}) = 1 \quad \text{true}$$

the system of equations $G_{\Sigma_1}(\vec{k}^*) = 0, G_{\Sigma_2}(\vec{k}^*) = 0$
has ∞ solutions but

the system $G_{\Sigma_1}(\vec{k}^*) = 0, G_{\Sigma_2}(\vec{k}^*) = 0, G_{\Sigma_3}(\vec{k}^*) = 0$
has NO solutions ...

$$\Rightarrow \vec{k}^* = (0.6223, 0.2953, 0)$$

solves $G_{\Sigma_1}(\vec{k}^*) = 0, G_{\Sigma_2}(\vec{k}^*) = 0$ and minimizes $|G_{\Sigma_3}(\vec{k}^*)|$

bcc

$\{\vec{R}\}$

$$R^2 \# \sum_R e^{+iG_r} = G_{\Sigma}^{(k)}$$

$$\sum_0 (000) \quad 0 \quad 1 \quad 1$$

$$\sum_1 (\pm\frac{1}{2}, \pm\frac{1}{2}, \pm\frac{1}{2}) \quad \frac{3}{4} \quad 8 \quad \cos(\pi k_x) \cos(\pi k_y) \cos(\pi k_z) \cdot 8$$

$$\sum_2 (\pm 1, 0, 0) \quad 1 \quad 6 \quad [\cos(2\pi k_x) + \cos(2\pi k_y) + \cos(2\pi k_z)] \cdot 2$$

$$\sum_3 (\pm 1, \pm 1, 0) \quad 2 \quad 12 \quad [\cos(2\pi k_x) \cos(2\pi k_y) + \cos(2\pi k_x) \cdot \cos(2\pi k_z) + \cos(2\pi k_y) \cdot \cos(2\pi k_z)] \cdot 4$$

⋮
⋮

$$G_{\Sigma_0}(k^*) = 1 \neq 0$$

the system $G_{\Sigma_1}(k^*) = G_{\Sigma_2}(k^*) = 0$ has ∞ solutions

the system $G_{\Sigma_1}(k^*) = G_{\Sigma_2}(k^*) = G_{\Sigma_3}(k^*) = 0$ has NO solutions

$$\Rightarrow k^* = (\frac{1}{6}, \frac{1}{6}, \frac{1}{2})$$

solves $G_{\Sigma_1}(k^*) = G_{\Sigma_2}(k^*) = 0$ and minimizes $|G_{\Sigma_3}(k^*)|$




FIG. 1. Mean-value point in the Brillouin zone of different cubic lattices. Solid circles are the mean-value points, whereas open circles indicate high-symmetry points.

how one can improve w.r.t. mean-value pt.?

fcc

	$G_{\Sigma_0}^{(k)}$	$G_{\Sigma_1}^{(k)}$	$G_{\Sigma_2}^{(k)}$	$G_{\Sigma_3}^{(k)}$
$\kappa^* = (0.6223, 0.2953, 0)$	1	0	0	$\neq 0$
$\kappa_1 = \left(\frac{1}{4}, \frac{1}{4}, \frac{1}{4}\right)$	1	6	0	0
$\kappa_2 = \left(\frac{1}{4}, \frac{1}{4}, \frac{3}{4}\right)$	1	-2	0	0
$\frac{1}{4}f(\kappa_1) + \frac{3}{4}f(\kappa_2)$	1	0	0	0

actually this is zero, up to Σ_7

$$\kappa_1 = \left(\frac{7}{8}, \frac{3}{8}, \frac{1}{8}\right), w_1 = \frac{6}{32}$$

$$\kappa_2 = \left(\frac{7}{8}, \frac{1}{8}, \frac{1}{8}\right), w_2 = \frac{3}{32}$$

$$\kappa_3 = \left(\frac{5}{8}, \frac{5}{8}, \frac{1}{8}\right), w_3 = \frac{3}{32}$$

$$\kappa_4 = \left(\frac{5}{8}, \frac{3}{8}, \frac{3}{8}\right), w_4 = \frac{3}{32}$$

$$\kappa_5 = \left(\frac{5}{8}, \frac{3}{8}, \frac{1}{8}\right), w_5 = \frac{6}{32}$$

$$\kappa_6 = \left(\frac{5}{8}, \frac{1}{8}, \frac{1}{8}\right), w_6 = \frac{3}{32}$$

$$\kappa_7 = \left(\frac{3}{8}, \frac{3}{8}, \frac{3}{8}\right), w_7 = \frac{1}{32}$$

$$\kappa_8 = \left(\frac{3}{8}, \frac{3}{8}, \frac{1}{8}\right), w_8 = \frac{3}{32}$$

$$\kappa_9 = \left(\frac{3}{8}, \frac{1}{8}, \frac{1}{8}\right), w_9 = \frac{3}{32}$$

$$\kappa_{10} = \left(\frac{1}{8}, \frac{1}{8}, \frac{1}{8}\right), w_{10} = \frac{1}{32}$$

$$\sum_{k=1}^{10} w_k = 1$$



$$\sum_{k=1}^{10} w_k G_{\Sigma_n}^{(k)} = 0$$

thn: $0 < n \leq 36$

$n=37$ gives $\Sigma \neq 0$

the weights are proportional to the number of κ vectors belonging to the same star

When performing calculations in Supercells (Defects, Alloys, Superlattices, Frozen Phonons) it is often convenient to use \mathbf{k} -points that are "equivalent" to those used in the bulk calculation in order to make the different energies exactly comparable.



If the symmetry of the crystal structure is lower than the symmetry of the Bravais lattice the number of inequivalent points increases and their weight changes

Special Points for superlattices

S.Y. Ren - J.D. Dow Phys. Rev. 38, 1999 (1988)

- A. Baldereschi Phys. Rev. B 7, 5212 (1973)
"Mean-Value Point in the Brillouin Zone"
- D.J. Chadi and M. Cohen
Phys. Rev. B 8, 5747 (1973)
"Special Points in the Brillouin Zone"
- H.J. Monkhorst and J.D. Pack
Phys. Rev. B 13, 5188 (1976)
"Special Points for Brillouin-zone integrations"
- D.J. Chadi and M. Cohen
Phys. Rev. B 16, 1746 (1977)
"Special Points for Brillouin zone integrations"
- J.D. Pack and H.J. Monkhorst
Phys. Rev. B 16, 1748 (1977)
"Special Points for BZ integrations"- a reply

see also

S.L. Cunningham, Phys. Rev B 10, 4988 (1974)
for special points in the hexagonal lattice.

Smearing techniques

$$n(r) = 2 \sum_v \frac{1}{(2\pi)^3} \int d\vec{k} \delta(\epsilon_F - \epsilon_v(\vec{k})) |\phi_{k,v}(r)|^2$$

↓ spin degeneracy
 we actually use a discrete grid

$$n(r) = 2 \sum_v \frac{1}{N_k} \sum_{\vec{k}_i} \delta(\epsilon_F - \epsilon_v(\vec{k}_i)) |\phi_{k_i,v}(r)|^2$$


the step function makes this integral badly convergent.

Possible solution: smooth the integral by substituting $\delta(\epsilon)$ with a smoother function



$$\delta(\epsilon)$$




$$\tilde{\delta}\left(\frac{\epsilon}{\sigma}\right)$$



$$\frac{d\delta(\epsilon)}{d\epsilon} = \delta'(\epsilon)$$



$$\frac{d\tilde{\delta}}{d\epsilon}\left(\frac{\epsilon}{\sigma}\right) = \frac{1}{\sigma} \tilde{\delta}'\left(\frac{\epsilon}{\sigma}\right)$$



The larger is σ the smoother is the function and therefore smaller the number of k-points in the grid needed to integrate accurately

Unfortunately

the larger is σ the more the function differs from the original one and the integral deviates from the $\sigma \rightarrow 0$ limit that corresponds to the physical situation.

Several smearing functions have been proposed

Fermi-Dirac


$$\mathcal{D}(x) = \frac{1}{1 + \exp(-x)}$$

$$\delta(x) = \frac{1}{2 + \exp(x) + \exp(-x)}$$

Gaussian: Fu and Ho, PRB 28, 5480 (1983)

$$\mathcal{D}(x) = A \cdot \text{erf}(\frac{x}{A})$$

$$\delta(x) = \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}}$$



Gaussian times polynomials

Methfessel and Paxton PRB 40 3616 (1989)
Marzari and Vanderbilt PRL 82 3296 (1999)

$$\mathcal{D}(x) = \int_{-\infty}^x \delta(y) dy$$

$$\delta(x) = \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}} p(k)$$

It can be shown that the "correct" energy is

$$E = 2 \sum_{k\sigma} \delta\left(\frac{\epsilon_F - \epsilon_{k\sigma}}{\sigma}\right) \langle \psi_{k\sigma} | -\frac{\hbar^2}{2m} \nabla^2 | \psi_{k\sigma} \rangle + \int V_0(r) n(r) d^3r \\ + \frac{e^2}{2} \int \frac{n(r) n(r')}{|r-r'|} d^3r d^3r' + E_{xc}[n] + \sigma \sum_{k\sigma} \delta_1 \left(\frac{\epsilon_F - \epsilon_{k\sigma}}{\sigma} \right)$$

for Fermi-Dirac smearing

$$\delta_1(x) = \int_{-\infty}^x y \frac{d}{dy} \left(\frac{1}{1+e^{-y}} \right) dy = \left[\frac{x}{1+e^{x}} \right] - \ln(1+e^x) \\ = \sigma \ln \sigma + (1-\sigma) \ln(1-\sigma)$$

that is the expression for the entropy for non-interacting electrons.

In fact in this case E is actually the Free Energy

since $\frac{dF}{dT} = -S$ we have (also for general smearing)

$$\frac{dE}{d\sigma} = \sum_{k\sigma} \delta_1 \left(\frac{\epsilon_F - \epsilon_{k\sigma}}{\sigma} \right) \approx \int_{-\infty}^{\infty} d\epsilon n(\epsilon) \int_{-\infty}^{\frac{(\epsilon_F - \epsilon)/\sigma}{\delta}} y d(y) dy \\ = - \sum_{k=0}^{\infty} \sigma^{2k+1} \frac{c_{k+1}}{(2k+1)!} \left. \frac{d^{2k} n}{d\epsilon^{2k}} \right|_{\epsilon_F}$$

$$c_k = \int_{-\infty}^{\infty} x^{2k} \delta(x) dx$$





FIG. 1. Total energy per atom of fcc-Pb crystal as a function of the smearing linewidth and for two different smearing functions: the Hermite-Gaussian function of order 1 (solid line) and the simple Gaussian function (dashed line).


HG_1 smearing function
 $\sigma=0.7 \text{ eV}$



smeared density of states
for Lead
 $\sigma=0.7 \text{ eV}$

DOS

$\varepsilon \text{ [eV]}$



SMEARING FUNCTIONS

- Fermi-Dirac

$$\tilde{\delta}(x) = \frac{1}{e^x + e^{-x} + 2}; \quad \tilde{\theta}(x) = \frac{1}{e^x + 1}$$

$$\sigma = K_B T; \quad T = 300K \implies \sigma = 0.026\text{eV}$$

- Gaussian

$$\tilde{\delta}(x) = \frac{1}{\sqrt{2\pi}} e^{-x^2/2}, \quad \tilde{\theta}(x) = \frac{1}{2} (1 + \text{erf}(x/\sqrt{2}))$$

- Hermite Polynomial \times Gaussian (HG_N) :
M.Methfessel and A.Paxton, **PRB 40**, 3616 (1989).

$$\tilde{\delta}(x) = \sum_{n=0}^N A_n H_{2n}(x) e^{-x^2}, \quad A_n = \frac{(-1)^n}{n! 4^n \sqrt{\pi}}$$

$$\tilde{\theta}(x) = \tilde{\theta}_0(x) - \sum_{n=1}^N A_n H_{2n-1}(x) e^{-x^2}$$

$$\text{with } \tilde{\theta}_0(x) = \frac{1}{2} (1 + \text{erf}(x))$$

THE ENERGY

Kohn-Sham (KS) kinetic energy functional

$$\begin{aligned}
 T_s[n] &= \int_{-\infty}^{\varepsilon_F} \varepsilon n(\varepsilon) d\varepsilon - \int V^{KS}(\mathbf{r}) n(\mathbf{r}) d\mathbf{r} \\
 &= \sum_i \left[-\frac{\hbar^2}{2m} \tilde{\theta}\left(\frac{\varepsilon_F - \varepsilon_i}{\sigma}\right) \langle \phi_i | \nabla^2 | \phi_i \rangle + \sigma \tilde{\theta}_1\left(\frac{\varepsilon_F - \varepsilon_i}{\sigma}\right) \right]
 \end{aligned}$$

$$\tilde{\theta}_1(x) = \int_{-\infty}^x y \delta(y) dy$$

Total Energy

$$E_{tot} = T_s[n] + E_{Hxc} + \int V^0(\mathbf{r}) n(\mathbf{r}) d\mathbf{r} + E_{ion}$$

KS equations


$$\left[-\frac{\hbar^2}{2m} \nabla^2 + V^0(\mathbf{r}) + V_{Hxc}(\mathbf{r}) - \varepsilon_i \right] \phi_i(\mathbf{r}) = 0$$

Hellmann-Feynman forces:

$$\frac{\partial E_{tot}}{\partial \mathbf{u}_s} = \int n(\mathbf{r}) \frac{\partial V^0(\mathbf{r})}{\partial \mathbf{u}_s} + \frac{\partial E_{ion}}{\partial \mathbf{u}_s}.$$

Pb:

TOTAL ENERGY vs SMEARING WIDTH



ENERGY DEPENDENCE ON SMEARING WIDTH

$$\begin{aligned}
 \frac{dE_{tot}}{d\sigma} &= \frac{\partial E_{tot}}{\partial \sigma} \\
 &= \sum_i \tilde{\theta}_1\left(\frac{\varepsilon_F - \varepsilon_i}{\sigma}\right) = \int_{-\infty}^{\infty} d\varepsilon \hat{n}_{\sigma}(\varepsilon) \int_{-\infty}^{\frac{\varepsilon_F - \varepsilon}{\sigma}} y \tilde{\delta}(y) dy \\
 &= - \sum_{k=0}^{\infty} \sigma^{2k+1} \frac{c_{k+1}}{(2k+1)!} \left. \frac{d^{2k} \hat{n}_{\sigma}}{d\varepsilon^{2k}} \right|_{\varepsilon_F} = - c_1 n(\varepsilon_F) \sigma - \frac{c_2 n''(\varepsilon_F)}{3!} \sigma^3 + \dots
 \end{aligned}$$

where $c_k = \int x^{2k} \tilde{\delta}(x) dx$

positive definite smearing functions


(Fermi-Dirac, Gaussian): $c_1 \neq 0$

$$E_{tot} \approx E_{tot}^{\sigma=0} - \frac{c_1}{2} n(\varepsilon_F) \sigma^2$$

non positive definite smearing functions, (HG1): $c_1 = 0$


$$E_{tot} \approx E_{tot}^{\sigma=0} - \frac{c_2}{4!} n''(\varepsilon_F) \sigma^4$$

Beryllium



Tetrahedra

given a regular grid of \vec{k} in the BZ
any small cube can be split in
6 tetrahedra that completely fill it



within each tetrahedron each energy band can be linearly interpolated

^{each of}
from the interpolated bands one can analytically calculate the fraction of the tetrahedron that corresponds to occupied states and then the weight of it

explicit formulas can be found in

Skriver, "the LMTO method" Springer-Verlag