On cubic Hodge integrals and random matrices

Boris Dubrovin, Di Yang

SISSA, via Bonomea 265, Trieste 34136, Italy

Abstract

A conjectural relationship between the GUE partition function with even couplings and certain special cubic Hodge integrals over the moduli spaces of stable algebraic curves is under consideration.

Contents

1	Introduction					
	1.1	Cubic Hodge partition function	1			
	1.2	GUE partition function with even couplings $\ldots \ldots \ldots$	3			
	1.3	From cubic Hodge integrals to random matrices. Main Conjecture	4			
	1.4	Computational aspects of the Main Conjecture: how do we verify it?	4			
	1.5	An explicit expression for F_g	7			
2	GUE free energy with even valencies					
	2.1	Calculating the GUE free energy from Frobenius manifold of \mathbb{P}^1 topological $\sigma\text{-model}$.	8			
	2.2	Proof of Prop. 1.4.1, Thm. 1.4.3	10			
3	Verification of the Main Conjecture for low genera					
	3.1	Genus 0	10			
	3.2	Genus 1,2	13			
	3.3	Genus 3,4	13			
\mathbf{A}	Exp	blicit formula for F_5	15			

1 Introduction

1.1 Cubic Hodge partition function

Let $\overline{\mathcal{M}}_{g,k}$ denote the Deligne–Mumford moduli space of stable curves of genus g with k distinct marked points. Denote by \mathcal{L}_i the i^{th} tautological line bundle over $\overline{\mathcal{M}}_{g,k}$, and $\mathbb{E}_{g,k}$ the rank g Hodge bundle.

Let $\psi_i := c_1(\mathcal{L}_i)$, $i = 1, \ldots, k$, and let $\lambda_i := c_i(\mathbb{E}_{g,k})$, $i = 0, \ldots, g$. Recall that the Hodge integrals over $\overline{\mathcal{M}}_{g,k}$, aka the intersection numbers of ψ - and λ -classes, are integrals of the form

$$\int_{\overline{\mathcal{M}}_{g,k}} \psi_1^{i_1} \cdots \psi_k^{i_k} \cdot \lambda_1^{j_1} \cdots \lambda_g^{j_g}, \qquad i_1, \dots, i_k, \, j_1, \dots, j_g \ge 0$$

Note that the dimension-degree matching implies that the above integrals vanish unless

$$3g - 3 + k = (i_1 + i_2 + \dots + i_k) + (j_1 + 2j_2 + 3j_3 + \dots + gj_g).$$

The particular case of *cubic Hodge integrals* of the form

$$\int_{\overline{\mathcal{M}}_{g,k}} \Lambda_g(p) \Lambda_g(q) \Lambda_g(r) \psi_1^{i_1} \cdots \psi_k^{i_k}, \qquad \frac{1}{p} + \frac{1}{q} + \frac{1}{r} = 0$$
(1.1.1)

was intensively studied after the formulation of the celebrated R. Gopakumar–M. Mariño–C. Vafa conjecture [17, 24] regarding the Chern–Simons/string duality. Here we denote

$$\Lambda_g(z) = \sum_{i=0}^g \lambda_i \, z^i$$

the Chern polynomial of $\mathbb{E}_{g,k}$. A remarkable expression for the cubic Hodge integrals of the form

$$\int_{\overline{\mathcal{M}}_{g,k}} \frac{\Lambda_g(p)\Lambda_g(q)\Lambda_g(r)}{(1-x_1\,\psi_1)\dots(1-x_k\,\psi_k)}, \quad k \ge 0$$

conjectured in [24] was proven in [18, 25]; for more about cubic Hodge integrals see in the subsequent papers [19, 20, 28, 12].

In the present paper we will deal with the specific case of Hodge integrals (1.1.1) with a pair of equal parameters among p, q, r; without loss of generality one can assume that p = q = -1, r = 1/2. So, the *special cubic Hodge integrals* of the form

$$\int_{\overline{\mathcal{M}}_{g,k}} \Lambda_g(-1) \Lambda_g(-1) \Lambda_g\left(\frac{1}{2}\right) \psi_1^{i_1} \cdots \psi_k^{i_k}$$
(1.1.2)

will be considered. Denote

$$\mathcal{H}(\mathbf{t};\epsilon) = \sum_{g\geq 0} \epsilon^{2g-2} \sum_{k\geq 0} \frac{1}{k!} \sum_{i_1,\dots,i_k\geq 0} t_{i_1}\cdots t_{i_k} \int_{\overline{\mathcal{M}}_{g,k}} \Lambda_g(-1) \Lambda_g(-1) \Lambda_g\left(\frac{1}{2}\right) \psi_1^{i_1}\cdots \psi_k^{i_k}$$
(1.1.3)

the generating function of these integrals. Here and below $\mathbf{t} = (t_0, t_1, ...)$ are independent variables, ϵ is a parameter. The exponential $e^{\mathcal{H}} =: Z_{\mathbb{E}}$ is called the cubic Hodge partition function while $\mathcal{H}(\mathbf{t}; \epsilon)$ is the cubic Hodge free energy. It can be written in the form of genus expansion

$$\mathcal{H}(\mathbf{t};\epsilon) = \sum_{g \ge 0} \epsilon^{2g-2} \mathcal{H}_g(\mathbf{t}) \tag{1.1.4}$$

where $\mathcal{H}_g(\mathbf{t})$ is called the genus g part of the cubic Hodge free energy, $g \ge 0$. Clearly $\mathcal{H}_0(\mathbf{t})$ coincides with the Witten–Kontsevich generating function of genus zero intersection numbers of ψ -classes

$$\mathcal{H}_{0}(\mathbf{t}) = \sum_{k \ge 0} \frac{1}{k!} \sum_{i_{1}, \dots, i_{k} \ge 0} t_{i_{1}} \cdots t_{i_{k}} \int_{\overline{\mathcal{M}}_{0,k}} \psi_{1}^{i_{1}} \cdots \psi_{k}^{i_{k}} = \sum_{k \ge 3} \frac{1}{k(k-1)(k-2)} \sum_{i_{1}+\dots+i_{k}=k-3} \frac{t_{i_{1}}}{i_{1}!} \cdots \frac{t_{i_{k}}}{i_{k}!}.$$
(1.1.5)

We note that an efficient algorithm for computing $\mathcal{H}_{g}(\mathbf{t}), g \geq 1$ was recently proposed in [12].

1.2 GUE partition function with even couplings

Let $\mathcal{H}(N)$ denote the space of $N \times N$ Hermitean matrices. Denote

$$dM = \prod_{i=1}^{N} dM_{ii} \prod_{i < j} d\text{Re}M_{ij} \, d\text{Im}M_{ij}$$

the standard unitary invariant volume element on $\mathcal{H}(N)$. The most studied Hermitean random matrix model is governed by the following GUE partition function with even couplings

$$Z_N(\mathbf{s}) = \frac{(2\pi)^{-N}}{Vol(N)} \int_{\mathcal{H}(N)} e^{-N \operatorname{tr} V(M;\mathbf{s})} dM.$$
(1.2.1)

Here, $V(M; \mathbf{s})$ is an **even** polynomial of M

$$V(M;\mathbf{s}) = \frac{1}{2}M^2 - \sum_{j\ge 1} s_j M^{2j},$$
(1.2.2)

or, more generally, a power series, by $\mathbf{s} = (s_1, s_2, s_3, ...)$ we denote the collection of coefficients¹ of V(M), and by Vol(N) the volume of the quotient of the unitary group over the maximal torus $[U(1)]^N$

$$Vol(N) = Vol\left(U(N)/[U(1)]^N\right) = \frac{\pi^{\frac{N(N-1)}{2}}}{G(N+1)}, \qquad G(N+1) = \prod_{n=1}^{N-1} n!.$$
(1.2.3)

The integral will be considered as a formal saddle point expansion with respect to the small parameter ϵ . Introduce the 't Hooft coupling parameter x by

 $x := N \epsilon.$

Reexpanding the free energy $\mathcal{F}_N(\mathbf{s}) := \log Z_N(\mathbf{s})$ in powers of ϵ and replacing the Barnes G-function by its asymptotic expansion yields²

$$\mathcal{F}(x,\mathbf{s};\epsilon) := \mathcal{F}_N(\mathbf{s})|_{N=\frac{x}{\epsilon}} - \frac{1}{12}\log\epsilon = \sum_{g\geq 0} \epsilon^{2g-2} \mathcal{F}_g(x,\mathbf{s}).$$
(1.2.4)

The GUE free energy $\mathcal{F}(x, \mathbf{s}; \epsilon)$ can be represented [21, 22, 1, 23] in the form

$$\mathcal{F}(x,\mathbf{s};\epsilon) = \frac{x^2}{2\epsilon^2} \left(\log x - \frac{3}{2}\right) - \frac{1}{12}\log x + \zeta'(-1) + \sum_{g\geq 2} \epsilon^{2g-2} \frac{B_{2g}}{4g(g-1)x^{2g-2}} + \sum_{g\geq 0} \epsilon^{2g-2} \sum_{k\geq 0} \sum_{i_1,\dots,i_k\geq 1} a_g(i_1,\dots,i_k) s_{i_1}\dots s_{i_k} x^{2-2g-(k-|i|)},$$
(1.2.5)

$$a_g(i_1, \dots, i_k) = \sum_{\Gamma} \frac{1}{\# \operatorname{Sym} \Gamma}$$
(1.2.6)

where the last summation is taken over all connected oriented ribbon graphs Γ of genus g with k vertices of valencies $2i_1, \ldots, 2i_k, \# \operatorname{Sym} \Gamma$ is the order of the symmetry group of Γ , and $|i| := i_1 + \cdots + i_k$.

Our goal is to compare the expansions (1.1.3) and (1.2.5).

¹The notation here is slightly different from that of [7, 8] where the coefficient of M^{2j} was denoted by s_{2j} .

²It is often called 1/N-expansion as $\epsilon \sim 1/N$.

1.3 From cubic Hodge integrals to random matrices. Main Conjecture.

It was already observed by E. Witten [27] that the GUE partition function with an even polynomial V(M) is tau-function of a particular solution to the Volterra (also called the discrete KdV) hierarchy. Recall that the first equation of the hierarchy (the Volterra lattice equation) reads

$$\dot{w}_n = w_n \left(w_{n+1} - w_{n-1} \right)$$

where

$$w_n = \frac{Z_{n+1}Z_{n-1}}{Z_n^2},$$

the time derivative is with respect to the variable $t = N s_1$. Other couplings s_k are identified with the time variables of higher flows of the hierarchy. On another side, the study [12] of integrable systems associated with the Hodge integrals³ suggested the following conjectural statement: the Hodge partition function $Z_{\mathbb{E}} = e^{\mathcal{H}}$ of the form (1.1.3) as function of independent parameters t_i is also a taufunction of the Volterra hierarchy. This observation provides a motivation for the main conjecture of the present paper.

It will be convenient to change normalisation of the GUE couplings. Put

$$\bar{s}_k := \left(\begin{array}{c} 2k\\k \end{array}\right) s_k.$$

Conjecture 1.3.1 (Main Conjecture) The following formula holds true

$$\sum_{g=0}^{\infty} \epsilon^{2g-2} \mathcal{F}_g(x, \mathbf{s}) + \epsilon^{-2} \left(-\frac{1}{2} \sum_{k_1, k_2 \ge 1} \frac{k_1 k_2}{k_1 + k_2} \bar{s}_{k_1} \bar{s}_{k_2} + \sum_{k \ge 1} \frac{k}{1+k} \bar{s}_k - x \sum_{k \ge 1} \bar{s}_k - \frac{1}{4} + x \right)$$
$$= \cosh\left(\frac{\epsilon \partial_x}{2}\right) \left[\sum_{g=0}^{\infty} \epsilon^{2g-2} 2^g \mathcal{H}_g\left(\mathbf{t}(x, \mathbf{s})\right) \right].$$
(1.3.1)

where

$$t_i(x, \mathbf{s}) := \sum_{k \ge 1} k^{i+1} \bar{s}_k - 1 + \delta_{i,1} + x \cdot \delta_{i,0}, \qquad i \ge 0.$$
(1.3.2)

1.4 Computational aspects of the Main Conjecture: how do we verify it?

We will check validity of the Main Conjecture for small genera. Begin with g = 0. Let us start with $\mathcal{H}_0(\mathbf{t})$. Instead of the explicit expansion (1.1.5) we use the following well known representation

$$\mathcal{H}_0 = \frac{v^3}{6} - \sum_{i \ge 0} t_i \frac{v^{i+2}}{i!(i+2)} + \frac{1}{2} \sum_{i,j \ge 0} t_i t_j \frac{v^{i+j+1}}{(i+j+1)\,i!\,j!}$$
(1.4.1)

where $v = v(t) = t_0 + \dots$ is the unique series solution to the equation

$$v = \sum_{i \ge 0} t_i \frac{v^i}{i!}.$$
 (1.4.2)

 $^{^{3}}$ The first example of an integrable system associated with *linear* Hodge integrals was investigated by A. Buryak. In this case the integrable system was proved to be Miura equivalent to the Intermediate Long Wave equation [3].

Here we recall that

$$v = \frac{\partial^2 \mathcal{H}_0(\mathbf{t})}{\partial t_0^2} = \sum_{k=1}^{\infty} \frac{1}{k} \sum_{i_1 + \dots + i_k = k-1} \frac{t_{i_1}}{i_1!} \dots \frac{t_{i_k}}{i_k!}$$
(1.4.3)

is a particular solution to the Riemann–Hopf hierarchy

$$\frac{\partial v}{\partial t_k} = \frac{v^k}{k!} \frac{\partial v}{\partial t_0}, \qquad k = 0, \, 1, \, 2, \dots.$$

For the genus zero GUE free energy $\mathcal{F}_0 = \mathcal{F}_0(x, \mathbf{s})$ one has a similar representation. Like above, introduce

$$u(x,\mathbf{s}) = \frac{\partial^2 \mathcal{F}_0(x,\mathbf{s})}{\partial x^2} \tag{1.4.4}$$

and put

$$w(x,\mathbf{s}) = e^{u(x,\mathbf{s})}.\tag{1.4.5}$$

Proposition 1.4.1 The function $w = w(x, \mathbf{s})$ is the unique series solution to the equation

$$w = x + \sum_{k \ge 1} k \, \bar{s}_k \, w^k, \qquad \bar{s}_k := \begin{pmatrix} 2k \\ k \end{pmatrix} s_k, \quad w(x, \mathbf{s}) = x + \dots$$
(1.4.6)

The genus zero GUE free energy \mathcal{F}_0 with even couplings has the following expression

$$\mathcal{F}_{0} = \frac{w^{2}}{4} - xw + \sum_{k \ge 1} \bar{s}_{k} \left(xw^{k} - \frac{k}{k+1}w^{k+1} \right) + \frac{1}{2} \sum_{k_{1},k_{2} \ge 1} \frac{k_{1}k_{2}}{k_{1}+k_{2}} \bar{s}_{k_{1}} \bar{s}_{k_{2}}w^{k_{1}+k_{2}} + \frac{x^{2}}{2} \log w. \quad (1.4.7)$$

Clearly w also satisfies the Riemann–Hopf hierarchy in a different normalization

$$\frac{\partial w}{\partial \bar{s}_k} = k \, w^k \frac{\partial w}{\partial x}, \qquad k \ge 1.$$

The solution can be written explicitly in the form essentially equivalent to (1.4.3)

$$w = \sum_{n=1}^{\infty} \frac{1}{n} \sum_{i_1 + \dots + i_n = n-1} \operatorname{wt}(i_1) \dots \operatorname{wt}(i_n) \bar{s}_{i_1} \dots \bar{s}_{i_n}$$

where we put $\bar{s}_0 = x$ and denote

$$\mathrm{wt}(i) = \left\{ \begin{array}{ll} 1, & i = 0 \\ \\ i, & \mathrm{otherwise.} \end{array} \right.$$

It is now straightforward to verify that the substitution (1.3.2) yields

$$e^{v(\mathbf{t}(x,\mathbf{s}))} = w(x,\mathbf{s}),$$
 i.e. $v(\mathbf{t}(x,\mathbf{s})) = u(x,\mathbf{s})$ (1.4.8)

and

$$\mathcal{H}_0\left(\mathbf{t}(x,\mathbf{s})\right) = \mathcal{F}_0\left(x,\mathbf{s}\right) - \frac{1}{2} \sum_{k_1,k_2 \ge 1} \frac{k_1 \, k_2}{k_1 + k_2} \, \bar{s}_{k_1} \, \bar{s}_{k_2} + \sum_{k \ge 1} \frac{k}{1+k} \, \bar{s}_k - x \, \sum_{k \ge 1} \, \bar{s}_k - \frac{1}{4} + x. \tag{1.4.9}$$

See in Sect. 2 for the details of this computation.

In order to proceed to higher genera we will use the method that goes back to the paper [5] by R. Dijkgraaf and E. Witten. The idea of this method is to express the positive genus free energy terms via the genus zero. Let us first explain this method for the Hodge free energy.

Theorem 1.4.2 ([12]) There exist functions $H_g(v, v_1, v_2, \ldots, v_{3g-2})$, $g \ge 1$ of independent variables v, v_1, v_2, \ldots such that

$$\mathcal{H}_{g}(\mathbf{t}) = H_{g}\left(v(\mathbf{t}), \frac{\partial v(\mathbf{t})}{\partial t_{0}}, \dots, \frac{\partial^{3g-2}v(\mathbf{t})}{\partial t_{0}^{3g-2}}\right), \quad g \ge 1.$$
(1.4.10)

Here $v(\mathbf{t})$ is given by eq. (1.4.3). Moreover, for any $g \ge 2$ the function H_g is a polynomial in the variables v_2, \ldots, v_{3g-2} with coefficients in $\mathbb{Q}\left[v_1, v_1^{-1}\right]$ (independent of v).

Explicitly,

$$H_1(v, v_1) = -\frac{1}{16}v + \frac{1}{24}\log v_1 \tag{1.4.11}$$

$$H_2(v_1, v_2, v_3, v_4) = \frac{7v_2}{2560} - \frac{v_1^2}{11520} + \frac{v_4}{1152v_1^2} - \frac{v_3}{320v_1} + \frac{v_2^3}{360v_1^4} + \frac{11v_2^2}{3840v_1^2} - \frac{7v_3v_2}{1920v_1^3}, \quad (1.4.12)$$

etc. The algorithm for computing the functions H_g can be found in [12]. They were used in the construction of the associated integrable hierarchy via the quasi-triviality transformation approach [10].

Let us now proceed to the higher genus terms for the random matrix free energy (recall that only even couplings are allowed).

Theorem 1.4.3 There exist functions $F_g(v, v_1, \ldots, v_{3g-2})$, $g \ge 1$ of independent variables v, v_1, v_2, \ldots such that

$$\mathcal{F}_g(x,\mathbf{s}) = F_g\left(u(x,\mathbf{s}), \frac{\partial u(x,\mathbf{s})}{\partial x}, \dots, \frac{\partial^{3g-2}u(x,\mathbf{s})}{\partial x^{3g-2}}\right), \quad g \ge 1.$$
(1.4.13)

Here

$$u(x, \mathbf{s}) = \frac{\partial^2 \mathcal{F}_0(x, \mathbf{s})}{\partial x^2} = \log w(x, \mathbf{s}).$$

Recall that the function $w(x, \mathbf{s})$ is determined from eq. (1.4.6).

Explicitly

$$F_1(v, v_1) = \frac{1}{12} \log v_1 + \text{const}$$
(1.4.14)

with const= $\frac{i\pi}{24} + \zeta'(-1)$,

$$F_2(v_1, v_2, v_3, v_4) = -\frac{v_2}{480} - \frac{v_1^2}{2880} + \frac{v_4}{288v_1^2} - \frac{v_3}{480v_1} + \frac{v_2^3}{90v_1^4} + \frac{v_2^2}{960v_1^2} - \frac{7v_3v_2}{480v_1^3}$$
(1.4.15)

etc. For any $g \ge 2$ the function F_g is a polynomial in the variables v_2, \ldots, v_{3g-2} with coefficients in $\mathbb{Q}\left[v_1, v_1^{-1}\right]$.

Using the fact that $\partial_{t_0} = \partial_x$ (see Section 3.2 below) along with the standard expansion

$$\cosh\left(\frac{\epsilon \,\partial_x}{2}\right) = 1 + \sum_{n \ge 1} \frac{1}{(2n)!} \left(\frac{\epsilon}{2}\right)^{2n} \partial_x^{2n}$$

we recast the Main Conjecture for $g \ge 1$ into a sequence of the following relationships between the functions F_g and H_g

$$F_1 = 2H_1 + \frac{v}{8} + \text{const}$$
(1.4.16)

and, for $g \geq 2$

$$F_g(v_1,\ldots,v_{3g-2}) = \frac{v_{2g-2}}{2^{2g}(2g)!} + \frac{D_0^{2g-2}H_1(v;v_1)}{2^{2g-3}(2g-2)!} + \sum_{m=2}^g \frac{2^{3m-2g}}{(2g-2m)!} D_0^{2(g-m)} H_m(v_1,\ldots,v_{3m-2}) \quad (1.4.17)$$

where the operator D_0 is defined by

$$D_0 = v_1 \frac{\partial}{\partial v} + \sum_{k \ge 1} v_{k+1} \frac{\partial}{\partial v_k}$$

For example,

$$F_2(v_1, v_2, v_3, v_4) = 4H_2(v_1, v_2, v_3, v_4) + \frac{1}{4}D_0^2H_1 + \frac{1}{384}v_2.$$
(1.4.18)

Eqs. (1.4.16), (1.4.18) can be easily verified (see below). In order to verify validity of eqs. (1.4.17) for any $g \ge 2$ we write a conjectural explicit expression for the functions $F_g(v_1, \ldots, v_{3g-2})$ responsible for the genus g random matrix free energies. This will be done in the next subsection.

1.5 An explicit expression for F_g

We first recall some notations. \mathbb{Y} will denote the set of all partitions. For any partition $\lambda \in \mathbb{Y}$ denote by $\ell(\lambda)$ the *length* of λ , by $\lambda_1, \lambda_2, \ldots, \lambda_{\ell(\lambda)}$ the non-zero components, $|\lambda| = \lambda_1 + \cdots + \lambda_{\ell(\lambda)}$ the *weight*, and by $m_i(\lambda)$ the *multiplicity* of i in λ . Put $m(\lambda)! := \prod_{i \ge 1} m_i(\lambda)!$. The set of all partitions of weight k will be denoted by \mathbb{Y}_k . For an arbitrary sequence of variables v_1, v_2, \ldots , denote $v_{\lambda} = v_{\lambda_1} \cdots v_{\lambda_{\ell(\lambda)}}$.

Conjecture 1.5.1 For any $g \ge 2$, the genus g GUE free energy F_q has the following expression

$$F_{g}(v_{1},\ldots,v_{3g-2}) = \frac{v_{2g-2}}{2^{2g}(2g)!} + \frac{1}{2^{2g-3}(2g-2)!} D_{0}^{2g-2} \left(-\frac{1}{16}v + \frac{1}{24}\log v_{1} \right) + \sum_{m=2}^{g} \frac{2^{3m-2g}}{(2g-2m)!}$$

$$\sum_{k=0}^{3m-3} \sum_{\substack{k_{1}+k_{2}+k_{3}=k\\0\leq k_{1},k_{2},k_{3}\leq m}} \frac{(-1)^{k_{2}+k_{3}}}{2^{k_{1}}} \sum_{\rho,\mu\in\mathbb{Y}_{3m-3-k}} \frac{\langle\lambda_{k_{1}}\lambda_{k_{2}}\lambda_{k_{3}}\tau_{\rho+1}\rangle_{g}}{m(\rho)!} Q^{\rho\mu} D_{0}^{2g-2m} \left(\frac{v_{\mu+1}}{v_{1}^{\ell(\mu)+m-1-k}} \right)$$

$$(1.5.1)$$

where for a partition $\mu = (\mu_1, \ldots, \mu_\ell)$, $\mu + 1$ denotes the partition $(\mu_1 + 1, \ldots, \mu_\ell + 1)$, $Q^{\rho\mu}$ is the so-called Q-matrix defined by

$$Q^{\rho\mu} = (-1)^{\ell(\rho)} \sum_{\substack{\mu^1 \in \mathbb{Y}_{\lambda_1}, \dots, \mu^{\ell(\rho)} \in \mathbb{Y}_{\lambda_{\ell}(\rho)} \\ \cup_{q=1}^{\ell(\rho)} \mu^q = \mu}} \prod_{q=1}^{\ell(\rho)} \frac{(\rho_q + \ell(\mu^q))! (-1)^{\ell(\mu^q)}}{m(\mu^q)! \prod_{j=1}^{\infty} (j+1)!^{m_j(\mu^q)}}.$$

In this formula we have used the notation

$$\langle \lambda_{k_1} \lambda_{k_2} \lambda_{k_3} \tau_{\nu} \rangle_g := \int_{\overline{\mathcal{M}}_{g,\ell}} \lambda_{k_1} \lambda_{k_2} \lambda_{k_3} \psi_1^{\nu_1} \dots \psi_{\ell}^{\nu_{\ell}}, \qquad \forall \nu = (\nu_1, \dots, \nu_{\ell}) \in \mathbb{Y}.$$

Details about Q-matrix can be found in [9]. Conj. 1.5.1 indicates that the special cubic Hodge integrals (1.1.2) naturally appear in the expressions for the higher genus terms of GUE free energy.

Organization of the paper In Sect. 2 we review the approach of [10, 7] to the GUE free energy, and prove Prop. 1.4.1 and Thm. 1.4.3. In Sect. 3 we verify Conj. 1.3.1 and Conj. 1.5.1 up to the genus 2 approximation, and give explicit formulae of \mathcal{F}_g for g = 3, 4, 5.

Acknowledgements We wish to thank Si-Qi Liu and Youjin Zhang for helpful discussions.

2 GUE free energy with even valencies

2.1 Calculating the GUE free energy from Frobenius manifold of \mathbb{P}^1 topological σ -model

It is known that the GUE partition function Z_N (with even and odd couplings) is the tau-function of a particular solution to the Toda lattice hierarchy. Using this fact, one of the authors in [7] developed an efficient algorithm of calculating of GUE free energy, which is an application of the general approach of [10, 6] for the particular example of the two-dimensional Frobenius manifold with potential

$$F = \frac{1}{2}u\,v^2 + e^u.$$

(Warning: only in this section, the notation v is different from that of the Introduction.) In this section, we give a brief reminder of this approach referring the readers to [10, 11, 7] for details.

Introduce two analytic functions $\theta_1(u, v; z), \theta_2(u, v; z)$ as follows

$$\theta_1(u,v;z) = -2e^{zv} \sum_{m=0}^{\infty} \left(-\frac{1}{2}u + c_m\right) e^{mu} \frac{z^{2m}}{m!^2} =: \sum_{p\ge 0} \theta_{1,p}(u,v) z^p$$
(2.1.1)

$$\theta_2(u,v;z) = z^{-1} \left(\sum_{m \ge 0} e^{mu + zv} \frac{z^{2m}}{(m!)^2} - 1 \right) =: \sum_{p \ge 0} \theta_{2,p}(u,v) \, z^p.$$
(2.1.2)

Here $c_m = \sum_{k=1}^m \frac{1}{k}$ denotes the *m*-th harmonic number.

Note that, as in the Introduction, we will only consider the GUE partition function with even couplings. The corresponding Euler–Lagrange equation [7, 6, 10] reads

$$x - w + \sum_{k \ge 1} (2k)! s_k \sum_{m=1}^k m \, w^m \, \frac{v^{2k-2m}}{(2k-2m)! \, m!^2} = 0 \tag{2.1.3}$$

$$-v + \sum_{k \ge 1} (2k)! s_k \sum_{m=0}^{k-1} w^m \frac{v^{2k-1-2m}}{(2k-1-2m)! m!^2} = 0$$
(2.1.4)

where $w = e^u$ (as in the Introduction). Note that we are only interested in the unique series solution $(v(x, \mathbf{s}), w(x, \mathbf{s}))$ of (2.1.3), (2.1.4) such that $v(x, \mathbf{0}) = 0$, $w(x, \mathbf{0}) = x$. It is then easy to see from eq. (2.1.4) that

$$v = v(x, \mathbf{s}) \equiv 0.$$

And eq. (2.1.3) becomes

$$x - w + \sum_{m \ge 1} s_{2m} m \, w^m \, \frac{(2m)!}{m!^2} = 0.$$
(2.1.5)

Define a family of analytic functions $\Omega_{\alpha,p;\beta,q}(u,v)$ by the following generating formula

$$\sum_{p,q\geq 0} \Omega_{\alpha,p;\beta,q} z^p y^q = \frac{1}{z+y} \left[\frac{\partial \theta_\alpha(z)}{\partial v} \frac{\partial \theta_\beta(y)}{\partial u} + \frac{\partial \theta_\alpha(z)}{\partial u} \frac{\partial \theta_\beta(y)}{\partial v} - \delta_{\alpha+\beta,3} \right], \qquad \alpha,\beta = 1,2.$$
(2.1.6)

The genus zero GUE free energy $\mathcal{F}_0(x, \mathbf{s})$ then has the following expression

$$\mathcal{F}_{0} = \frac{1}{2} \sum_{p,q \ge 2} (2p)! (2q)! \, s_{p} s_{q} \, \Omega_{2,2p-1;2,2q-1} + x \sum_{q \ge 1} (2q)! \, s_{q} \, \Omega_{1,0;2,2q-1} - x \, \Omega_{1,0;2,1} \\ + \frac{1}{2} (1 - 2s_{1})^{2} \, \Omega_{2,1;2,1} + \sum_{q \ge 2} (2s_{1} - 1) \, (2q)! \, s_{q} \, \Omega_{2,1;2,2q-1} + \frac{1}{2} x^{2} \, \Omega_{1,0;1,0}.$$
(2.1.7)

The higher genus terms in the 1/N expansion of the GUE free energy can be determined recursively from the *loop equation* [10, 7] for a sequence of functions

$$F_g = F_g(u, v, u_1, v_1, \dots, v_{3g-2}, u_{3g-2}), \quad g \ge 1.$$

This equation has the following form

$$\sum_{r\geq 0} \left[\frac{\partial\Delta\mathcal{F}}{\partial v_r} \left(\frac{v-\lambda}{D} \right)_r - 2 \frac{\partial\Delta\mathcal{F}}{\partial u_r} \left(\frac{1}{D} \right)_r \right] \\ + \sum_{r\geq 1} \sum_{k=1}^r \binom{r}{k} \left(\frac{1}{\sqrt{D}} \right)_{k-1} \left[\frac{\partial\Delta\mathcal{F}}{\partial v_r} \left(\frac{v-\lambda}{\sqrt{D}} \right)_{r-k+1} - 2 \frac{\partial\Delta\mathcal{F}}{\partial u_r} \left(\frac{1}{\sqrt{D}} \right)_{r-k+1} \right] \\ = D^{-3} e^u \left(4 e^u + (v-\lambda)^2 \right) - \epsilon^2 \sum_{k,l} \left[\frac{1}{4} S(\Delta\mathcal{F}, v_k, v_l) \left(\frac{v-\lambda}{\sqrt{D}} \right)_{k+1} \left(\frac{v-\lambda}{\sqrt{D}} \right)_{l+1} \right] \\ - S(\Delta\mathcal{F}, v_k, u_l) \left(\frac{v-\lambda}{\sqrt{D}} \right)_{k+1} \left(\frac{1}{\sqrt{D}} \right)_{l+1} + S(\Delta\mathcal{F}, u_k, u_l) \left(\frac{1}{\sqrt{D}} \right)_{k+1} \left(\frac{1}{\sqrt{D}} \right)_{l+1} \right] \\ - \frac{\epsilon^2}{2} \sum_k \left[\frac{\partial\Delta\mathcal{F}}{\partial v_k} \frac{4 e^u (v-\lambda) u_l - T v_l}{D^3} + \frac{\partial\Delta\mathcal{F}}{\partial u_k} \frac{4 (v-\lambda) v_l - T u_l}{D^3} \right] (e^u)_{k+1}$$
(2.1.8)

where $\Delta \mathcal{F} = \sum_{g \ge 1} \epsilon^{2g} F_g$, $D = (v - \lambda)^2 - 4e^u$, $T = (v - \lambda)^2 + 4e^u$, $S(f, a, b) := \frac{\partial^2 f}{\partial a \partial b} + \frac{\partial f}{\partial a} \frac{\partial f}{\partial b}$, and f_r stands for $\partial_x^r(f)$. Solution $\Delta \mathcal{F}$ of (2.1.8) exists and is unique up to an additive constant. F_g is a polynomial in $u_2, v_2, \ldots, u_{3g-2}, v_{3g-2}$. For $g \ge 2$, F_g is a rational function of u_1, v_1 . Then [10] the genus g term in the expansion (1.2.4), in the particular case of even couplings only, reads

$$\mathcal{F}_g(x,\mathbf{s}) = F_g\left(u(x,\mathbf{s}), v = 0, \frac{\partial u(x,\mathbf{s})}{\partial x}, v_x = 0, \dots, \frac{\partial^{3g-2}u(x,\mathbf{s})}{\partial x^{3g-2}}, v_{3g-2} = 0\right), \qquad g \ge 1$$

This procedure will be used in the next subsection.

2.2 Proof of Prop. 1.4.1, Thm. 1.4.3

Proof of Prop. 1.4.1. Noting that

$$\theta_2(u,0,z) = z^{-1} \left(\sum_{m \ge 0} w^m \frac{z^{2m}}{(m!)^2} - 1 \right),$$

$$\partial_v \theta_2(u,0,z) = \sum_{m \ge 0} w^m \frac{z^{2m}}{(m!)^2}, \qquad \partial_u \theta_2(u,0,z) = \sum_{m \ge 0} m w^m \frac{z^{2m-1}}{(m!)^2}$$

and using (2.1.6) we have

$$\sum_{p,q\geq 0} \Omega_{2,p;2,q} z^p y^q = \frac{\sum_{m\geq 0} w^m \frac{z^{2m}}{(m!)^2} \sum_{m\geq 0} mw^m \frac{y^{2m-1}}{(m!)^2} + \sum_{m\geq 0} mw^m \frac{z^{2m-1}}{(m!)^2} \sum_{m\geq 0} w^m \frac{y^{2m}}{(m!)^2}}{z+y}.$$

It follows that if p + q is odd then $\Omega_{2,p;2,q}$ vanishes; otherwise, we have

$$\Omega_{2,p;2,q} = \frac{w^{\frac{p+q}{2}+1}}{\left(1+\frac{p+q}{2}\right) \left[\left(\frac{p}{2}\right)!\right]^2 \left[\left(\frac{q}{2}\right)!\right]^2}, \qquad p,q \text{ are both even};$$
(2.2.1)
$$\frac{p+1}{q+1} w^{\frac{p+q}{2}+1}$$

$$\Omega_{2,p;2,q} = \frac{\frac{p+1}{2}\frac{q+1}{2}w^{-2}+1}{\left(1+\frac{p+q}{2}\right)\left[\left(\frac{p+1}{2}\right)!\right]^2\left[\left(\frac{q+1}{2}\right)!\right]^2}, \quad p,q \text{ are both odd.}$$
(2.2.2)

Substituting these expressions in (2.1.7) we obtain

$$\mathcal{F}_{0} = \frac{1}{2}x^{2}u + \frac{1}{2}\sum_{k_{1},k_{2}\geq0}(2k_{1}+2)!(2k_{2}+2)!s_{k_{1}+1}s_{k_{2}+1}\frac{(k_{1}+1)(k_{2}+1)w^{k_{1}+k_{2}+2}}{(k_{1}+k_{2}+2)\left[(k_{1}+1)!\right]^{2}\left[(k_{2}+1)!\right]^{2}} + x\sum_{k\geq0}(2k+2)!s_{k+1}\frac{w^{k+1}}{(k+1)!^{2}} - xw + (1-4s_{1})\frac{w^{2}}{4} - \sum_{k\geq1}(2k+2)!s_{k+1}\frac{(k+1)w^{k+2}}{(k+2)\left[(k+1)!\right]^{2}}.$$

Equation (1.4.6) is already proved in (2.1.5). The proposition is proved.

Proof of Theorem 1.4.3. For g = 1, 2, taking $v = v_1 = v_2 = \cdots = 0$ in the general expressions of $F_g(u, v, u_1, v_1, \ldots, u_{3g-2}, v_{3g-2})$ [10, 11] one obtains (1.4.14) and (1.4.15). For any $g \ge 1$, the existence of $F_g(u, u_1, \ldots, u_{3g-2})$ such that

$$\mathcal{F}_g(x,\mathbf{s}) = F_g\left(u(x,\mathbf{s}), \frac{\partial u(x,\mathbf{s})}{\partial x}, \dots, \frac{\partial^{3g-2}u(x,\mathbf{s})}{\partial x^{3g-2}}\right)$$

is a direct result of [10, 11] when taking $v = v_1 = v_2 = \cdots = 0$ in $F_g(u, v, u_1, v_1, \dots, u_{3g-2}, v_{3g-2})$.

3 Verification of the Main Conjecture for low genera

3.1 Genus 0

Recall that the genus zero cubic Hodge free energy can be expressed as

$$\mathcal{H}_0(\mathbf{t}) = \frac{1}{2} \sum_{i,j \ge 0} \tilde{t}_i \, \tilde{t}_j \, \Omega_{i;j}(v(\mathbf{t})).$$

where $\tilde{t}_i = t_i - \delta_{i,1}$, $\mathbf{t} = (t_0, t_1, t_2, \dots)$, $\Omega_{i;j}$ are polynomials in v given by

$$\Omega_{i;j}(v) = \frac{v^{i+j+1}}{(i+j+1)\,i!\,j!},$$

and $v(\mathbf{t})$ is the unique series solution to the following Euler–Lagrange equation of the one-dimensional Frobenius manifold

$$v = \sum_{i \ge 0} t_i \frac{v^i}{i!}.$$

(Warning: the above v is the flat coordinate of the one-dimensional Frobenius manifold; avoid confusing with v in Section 2 where (u, v) are flat coordinates of the two-dimensional Frobenius manifold of \mathbb{P}^1 topological σ -model.)

Let us consider the following substitution of time variables

$$t_i = \sum_{k \ge 1} k^{i+1} \bar{s}_k - 1 + \delta_{i,1} + x \cdot \delta_{i,0}, \qquad i \ge 0.$$

Note that with this substitution the cubic Hodge free energies will be considered to be expanded at x = 1. We have $\tilde{t}_i = \sum_{k \ge 1} k^{i+1} \bar{s}_k - 1 + x \cdot \delta_{i,0}$, and so

$$\mathcal{H}_{0} = \frac{1}{2} \sum_{i,j \geq 0} \tilde{t}_{i} \tilde{t}_{j} \Omega_{i;j}(v(\mathbf{t})) \\
= \frac{1}{2} \sum_{i,j \geq 0} \left(\sum_{k_{1} \geq 1} k_{1}^{i+1} \bar{s}_{k_{1}} - 1 + x \cdot \delta_{i,0} \right) \left(\sum_{k_{2} \geq 1} k_{2}^{j+1} \bar{s}_{k_{2}} - 1 + x \cdot \delta_{j,0} \right) \frac{v^{i+j+1}}{(i+j+1)\,i!\,j!} \\
= \frac{1}{2} \sum_{i,j \geq 0} \sum_{k_{1},k_{2} \geq 1} k_{1}^{i+1} k_{2}^{j+1} \bar{s}_{k_{1}} \bar{s}_{k_{2}} \frac{v^{i+j+1}}{(i+j+1)\,i!\,j!} - \sum_{i,j \geq 0} \sum_{k_{1} \geq 1} k_{1}^{i+1} \bar{s}_{k_{1}} \frac{v^{i+j+1}}{(i+j+1)\,i!\,j!} \\
+ x \sum_{i,j \geq 0} \sum_{k_{1} \geq 1} k_{1}^{i+1} \bar{s}_{k_{1}} \delta_{j,0} \frac{v^{i+j+1}}{(i+j+1)\,i!\,j!} + \frac{1}{2} \sum_{i,j \geq 0} \frac{v^{i+j+1}}{(i+j+1)\,i!\,j!} \\
- x \sum_{i,j \geq 0} \delta_{j,0} \frac{v^{i+j+1}}{(i+j+1)\,i!\,j!} + \frac{x^{2}}{2} \sum_{i,j \geq 0} \delta_{i,0} \delta_{j,0} \frac{v^{i+j+1}}{(i+j+1)\,i!\,j!}.$$
(3.1.1)

We simplify it term by term:

$$\begin{split} \frac{x^2}{2} \sum_{i,j \ge 0} \delta_{i,0} \, \delta_{j,0} \, \frac{v^{i+j+1}}{(i+j+1)\,i!\,j!} &= \frac{x^2}{2}v, \\ x \sum_{i,j \ge 0} \, \delta_{j,0} \, \frac{v^{i+j+1}}{(i+j+1)\,i!\,j!} &= x \, (e^v - 1), \\ \frac{1}{2} \sum_{i,j \ge 0} \frac{v^{i+j+1}}{(i+j+1)\,i!\,j!} &= \frac{1}{2} \sum_{\ell \ge 0} \sum_{i=0}^{\ell} \frac{v^{\ell+1}\ell!}{(\ell+1)!\,i!\,(\ell-i)!} &= \frac{1}{2} \sum_{\ell \ge 0} \frac{v^{\ell+1}\,2^{\ell}}{(\ell+1)!} &= \frac{1}{4} (e^{2v} - 1), \\ x \sum_{i,j \ge 0} \sum_{k_1 \ge 1} k_1^{i+1} \, \bar{s}_{k_1} \, \delta_{j,0} \, \frac{v^{i+j+1}}{(i+j+1)\,i!\,j!} &= x \sum_{i \ge 0} \sum_{k_1 \ge 1} k_1^{i+1} \, \bar{s}_{k_1} \, \frac{v^{i+1}}{(i+1)!} &= x \sum_{k \ge 1} \bar{s}_k \, (e^{kv} - 1), \end{split}$$

$$\sum_{i,j\geq 0} \sum_{k_1\geq 1} k_1^{i+1} \bar{s}_{k_1} \frac{v^{i+j+1}}{(i+j+1)\,i!\,j!} = \sum_{k\geq 1} k \,\bar{s}_k \sum_{\ell\geq 0} (1+k)^\ell \frac{v^{\ell+1}}{(\ell+1)!} = \sum_{k\geq 1} \frac{k}{1+k} \,\bar{s}_k \left(e^{(1+k)v} - 1\right),$$

$$\frac{1}{2} \sum_{i,j\geq 0} \sum_{k_1,k_2\geq 1} k_1^{i+1} \,k_2^{j+1} \,\bar{s}_{k_1} \,\bar{s}_{k_2} \frac{v^{i+j+1}}{(i+j+1)\,i!\,j!} = \frac{1}{2} \sum_{k_1,k_2\geq 1} k_1 \,k_2 \,\bar{s}_{k_1} \,\bar{s}_{k_2} \sum_{\ell\geq 0} \frac{v^{\ell+1}}{(\ell+1)!} (k_1+k_2)^\ell$$

$$= \frac{1}{2} \sum_{k_1,k_2\geq 1} \frac{k_1 \,k_2}{k_1+k_2} \,\bar{s}_{k_1} \,\bar{s}_{k_2} \left(e^{(k_1+k_2)v} - 1\right).$$

Let $w = e^v$. We have

$$\mathcal{H}_{0} = \frac{1}{2} \sum_{k_{1},k_{2}\geq 1} \frac{k_{1}k_{2}}{k_{1}+k_{2}} \bar{s}_{k_{1}} \bar{s}_{k_{2}} \left(w^{k_{1}+k_{2}}-1\right) - \sum_{k\geq 1} \frac{k}{1+k} \bar{s}_{k} \left(w^{1+k}-1\right) + x \sum_{k\geq 1} \bar{s}_{k} \left(w^{k}-1\right) + \frac{1}{4} (w^{2}-1) - x \left(w-1\right) + \frac{x^{2}}{2} \log w.$$

On the other hand, recall from Prop. 1.4.1 that the genus zero GUE free energy with even couplings has the form

$$\mathcal{F}_0 = \frac{w^2}{4} - x \, w + \sum_{k \ge 1} \bar{s}_k \left(x \, w^k - \frac{k}{k+1} w^{k+1} \right) + \frac{1}{2} \sum_{k_1, k_2 \ge 1} \frac{k_1 k_2}{k_1 + k_2} \bar{s}_{k_1} \bar{s}_{k_2} w^{k_1 + k_2} + \frac{x^2}{2} \log w.$$

Here w is the power series solution to

$$w = x + \sum_{k \ge 1} k \, \bar{s}_k \, w^k.$$

Recall that $w = e^u$; so

$$e^u = x + \sum_{k \ge 1} k \, \bar{s}_k \, e^{ku}.$$

Namely,

$$1 + \sum_{j \ge 1} \frac{u^j}{j!} = x + \sum_{k \ge 1} k \,\bar{s}_k \,\left(1 + \sum_{j \ge 1} \frac{k^j \, u^j}{j!} \right).$$

It follows that

$$u(x, \mathbf{s}) = v(\mathbf{t}(x, \mathbf{s})). \tag{3.1.2}$$

We conclude that

$$\mathcal{H}_0(\mathbf{t}(x,\mathbf{s})) - \mathcal{F}_0(x,\mathbf{s}) = -\frac{1}{2} \sum_{k_1,k_2 \ge 1} \frac{k_1 k_2}{k_1 + k_2} \bar{s}_{k_1} \bar{s}_{k_2} + \sum_{k \ge 1} \frac{k}{1+k} \bar{s}_k - x \sum_{k \ge 1} \bar{s}_k - \frac{1}{4} + x.$$
(3.1.3)

This finishes the proof of the genus zero part of the Main Conjecture.

3.2 Genus 1, 2

Note that the substitution (1.3.2)

$$(t_0, t_1, t_2, \dots) \mapsto (x, \overline{s}_1, \overline{s}_2, \dots)$$

satisfies that

$$\frac{\partial}{\partial x} = \frac{\partial}{\partial t_0},\tag{3.2.1}$$

$$\frac{\partial}{\partial \bar{s}_k} = \sum_{i \ge 0} k^{i+1} \frac{\partial}{\partial t_i}, \qquad k \ge 1.$$
(3.2.2)

In particular, we have

$$\frac{\partial v}{\partial t_0}(\mathbf{t}(x,\mathbf{s})) = \frac{\partial v(\mathbf{t}(x,\mathbf{s}))}{\partial x} = \frac{\partial u(x,\mathbf{s})}{\partial x}.$$

The last equality is due to (3.1.2).

Recall, from the algorithm of [12], that the genus 1 special cubic Hodge free energy is given by

$$H_1(v; v_1) = \frac{1}{24} \log v_1 - \frac{v}{16}.$$

 So

$$2H_1(v;v_1) + \frac{v}{8} + \frac{i\pi}{24} + \zeta'(-1) = \frac{1}{12}\log v_1 + \frac{i\pi}{24} + \zeta'(-1).$$

This proves the genus 1 part of the Main Conjecture.

The genus 2 term of the special cubic Hodge free energy is given by

$$H_2(v_1, v_2, v_3, v_4) = \frac{7v_2}{2560} - \frac{v_1^2}{11520} + \frac{v_4}{1152v_1^2} - \frac{v_3}{320v_1} + \frac{v_2^3}{360v_1^4} + \frac{11v_2^2}{3840v_1^2} - \frac{7v_3v_2}{1920v_1^3}$$

 So

$$4H_2 + \frac{1}{4}D_0^2H_1 + \frac{1}{384}v_2 = 4H_2(v_1, v_2, v_3, v_4) - \frac{5}{384}v_2 + \frac{1}{96}\left[\frac{v_3}{v_1} - \left(\frac{v_2}{v_1}\right)^2\right]$$
$$= -\frac{v_2}{480} - \frac{v_1^2}{2880} + \frac{v_4}{288v_1^2} - \frac{v_3}{480v_1} + \frac{v_2^3}{90v_1^4} + \frac{v_2^2}{960v_1^2} - \frac{7v_3v_2}{480v_1^3}$$
$$= F_2(v_1, v_2, v_3, v_4).$$

This proves the genus 2 part of the Main Conjecture.

3.3 Genus 3, 4

Using the Main Conjecture along with the algorithm of [12], we obtain the following two statements.

Conjecture 3.3.1 The genus 3 GUE free energy is given by

$F_3(u_1,\ldots,u_7)$	
$13u_4 u_2^2 u_1^4 u_7 u_6 u_5$	
$-\frac{1}{120960} + \frac{1}{24192} - \frac{1}{725760} + \frac{1}{10368u_1^3} - \frac{1}{5760u_1^2} - \frac{1}{13440u_1}$	
$103u_4^2$ $59u_3^3$ u_3^2 u_3u_1 $5u_2^6$ $13u_2^5$	
$-\frac{-1}{60480u_1^4} + \frac{1}{8064u_1^5} + \frac{1}{2688u_1^2} + \frac{1}{12096} - \frac{1}{81u_1^8} - \frac{1}{1890u_1^6}$	
$5u_2^4$ u_2^3 $7u_6u_2$ $53u_3u_5$ $353u_5u_2^2$ u_5u_2	
$+\frac{1}{5376u_1^4}-\frac{1}{9072u_1^2}-\frac{1}{5760u_1^4}-\frac{1}{20160u_1^4}+\frac{1}{40320u_1^5}+\frac{1}{840u_1^3}$	
$89u_3u_4$ $83u_4u_2^3$ $211u_4u_2^2$ u_4u_2 $59u_3u_2^4$	
$+\frac{1}{40320u_1^3} - \frac{1}{1890u_1^6} - \frac{1}{40320u_1^4} + \frac{1}{2016u_1^2} + \frac{1}{378u_1^7}$	
$1993u_3u_2^3 u_3u_2^2 83u_3^2u_2^2 19u_3u_2 17u_3^2u_2 1273u_3u_4u_2$	
$+\frac{1}{120960u_1^5} - \frac{1}{576u_1^3} - \frac{1}{896u_1^6} + \frac{1}{120960u_1} - \frac{1}{2240u_1^4} + \frac{1}{40320u_1^5}.$	
	(3.3.1)

Conjecture 3.3.2 The genus 4 GUE free energy is given by

$F_4(u_1,\ldots,u_{10})$
$=\frac{1852u_2^9}{12}+\frac{151u_2^8}{12}-\frac{101u_2^7}{12}-\frac{772u_3u_2^7}{12}$
$1215u_1^{12}$ $675u_1^{10}$ $12600u_1^8$ $135u_1^{11}$
$9904u_4u_2^6$ $1165u_2^6$ $2851u_3u_2^6$ $14903u_3^2u_2^5$ $70261u_3u_2^5$
$+ \frac{1}{6075u_1^{10}} - \frac{1}{1161216u_1^6} - \frac{1}{3600u_1^9} + \frac{1}{2160u_1^{10}} + \frac{1}{3225600u_1^7}$
$2573u_4u_2^5$ u_2^5 $2243u_5u_2^5$ $195677u_3^2u_2^4$ $3197u_3u_2^4$
$+ \frac{1}{10800u_1^8} + \frac{1}{7200u_1^4} - \frac{1}{6480u_1^9} + \frac{1}{230400u_1^8} + \frac{1}{967680u_1^5}$
$+\frac{12907u_6u_2^4}{10259u_4u_2^4} - \frac{10259u_4u_2^4}{10259u_4u_2^4} - \frac{22153u_5u_2^4}{101503u_3u_4u_2^4} - \frac{1023u_4u_2^4}{1823u_4^2u_2^3} - \frac{101503u_3u_4u_2^4}{1823u_4u_2^4} - \frac{10150u_4u_2^4}{1823u_4u_2^4} - \frac{10050u_4u_2^4}{1823u_4u_2^4} - \frac{10050u_4u_4u_2^4}{1823u_4u_2^4} - \frac{10050u_4u_4u_4^4}{1823u_4u_4u_4} - \frac{10050u_4u_4u_4}{1823u_4u_4} - \frac{10050u_4u_4u_4}{1823u_4u_4} - \frac{10050u_4u_4}{1823u_4u_4} - \frac{10050u_4u_4}{1823u_4u_4} - \frac{10050u_4u_4}{1823u_4} - \frac{10050u_4u_4}{1823u_4} - \frac{10050u_4u_4}{1823u_4} - \frac{10050u_4u_4}{1823u_4} - \frac{10050u_4}{1823u_4} - \frac{10050u_4u_4}{1823u_4} $
$+ \frac{1}{226800u_1^8} - \frac{1}{1935360u_1^6} - \frac{1}{414720u_1^7} - \frac{1}{32400u_1^9} + \frac{1}{5670u_1^8} - $
$+ \frac{415273u_3u_5u_2^3}{49u_3u_2^3} + \frac{97u_5u_2^3}{26879u_6u_2^3} + \frac{26879u_6u_2^3}{2} + \frac{u_2^3}{49u_3u_2^3} + \frac{49u_3u_2^3}{49u_3u_2^3} + \frac{49u_3u_2^3}{49u_3u_2^3} + 1000000000000000000000000000000000000$
$+ \frac{-829440u_1^8}{120960u_1^5} + \frac{-2903040u_1^6}{2903040u_1^6} + \frac{-7257600}{7257600} - \frac{-138240u_1^3}{138240u_1^3}$
$5137u_4u_2^3 \qquad 877u_3^2u_2^3 \qquad 812729u_3u_4u_2^3 \qquad 212267u_7u_2^3 \qquad 305129u_3^3u_2^3$
$-\frac{-1}{4354560u_1^4} - \frac{-1}{57600u_1^6} - \frac{-1}{2073600u_1^7} - \frac{-1}{29030400u_1^7} - \frac{-103680u_1^9}{103680u_1^9} - \frac{-103680u_1^9}{10000u_1^9} - \frac{-10000u_1^9}{10000u_1^9} - \frac{-10000u_1^9}{100000$
$u_1^2 u_2^2$ 1379 $u_4^2 u_2^2$ 13138507 $u_3^2 u_4 u_2^2$ 2417 $u_3 u_4 u_2^2$ 17 $u_4 u_2^2$
$+\frac{1}{460800} + \frac{1}{34560u_1^6} + \frac{1}{9676800u_1^8} + \frac{1}{537600u_1^5} + \frac{1}{138240u_1^2}$
$2143u_3u_5u_2^2 + 449u_5u_2^2 + 2323u_8u_2^2 - 2623u_3^2u_2^2 - 443u_6u_2^2$
$+ \frac{1}{34560u_1^6} + \frac{1}{1451520u_1^3} + \frac{1}{3225600u_1^6} - \frac{1}{967680u_1^4} - \frac{1}{9676800u_1^4} - \frac{1}{967$
$667u_7u_2^2 \qquad 192983u_3^3u_2^2 \qquad 60941u_3u_6u_2^2 \qquad 171343u_4u_5u_2^2 \qquad 22809u_3^4u_2$
$-\frac{-\frac{-1}{537600u_1^5}-\frac{-1}{691200u_1^7}-\frac{-1}{1075200u_1^7}-\frac{-1}{1935360u_1^7}+\frac{-1}{71680u_1^8}}{71680u_1^8}$
$1747u_3^3u_2$ $7u_3^2u_2$ $9221u_5^2u_2$ $17u_1u_3u_2$ $78533u_3^2u_4u_2$
$+\frac{1}{806400u_1^5}+\frac{1}{38400u_1^2}+\frac{1}{1935360u_1^6}+\frac{1}{3225600}+\frac{1}{691200u_1^6}$
$+\frac{18713u_3u_4u_2}{1000}+\frac{15179u_4u_6u_2}{1000}+\frac{20639u_3u_7u_2}{1000}+\frac{37u_8u_2}{1000}+\frac{1000}$
$14515200u_1^3 + 1935360u_1^6 + 4838400u_1^6 + 302400u_1^4 - 86400$
$11u_5u_2 \qquad 923u_6u_2 \qquad 113u_7u_2 \qquad 55u_4^2u_2 \qquad 419u_3u_5u_2$
$-\frac{-362880u_1}{362880u_1} - \frac{-14515200u_1^2}{14515200u_1^2} - \frac{-9676800u_1^3}{9676800u_1^3} - \frac{-387072u_1^4}{387072u_1^4} - \frac{-1935360u_1^4}{1935360u_1^4}$

$1411u_4u_5u_2$ 7	$u_9u_2 = 1751u_3u_6u_2$	$12035u_3^2u_5u_2$ 442	$01u_3u_4^2u_2$
$-\frac{138240u_1^5}{138}$	$\overline{8240u_1^5}$ 268800 u_1^5 -	$96768u_1^7$ 27	$76480u_1^7$
$1549u_3^4$ 93'	$7u_3^3$, $229u_4^3$, 19	$9u_5^2$ $u_1^3u_3$ 94	$9u_3u_4u_5$
$+\frac{115200u_1^6}{29030}$	$\overline{040u_1^3} + \overline{62208u_1^6} + \overline{460}$	$\overline{80u_1^4} + \overline{691200} + \overline{5}$	$5296u_1^6$
$59u_3^2u_6$ $73u_4^2$	$u_6 = 1777u_3u_7 = 1000$	$143u_7$ $31u_8$	u_{10}
$+\frac{10752u_1^6}{10752u_1^6}+\frac{107520}{107520}$	$\overline{0u_1^4} + \overline{4838400u_1^4} + \overline{145}$	$15200u_1 + 9676800u$	$\frac{1}{1}^{2} + \frac{1}{497664u_{1}^{4}}$
$u_3^2 u_1 u_5$	$73u_6 u_1^6$	$19u_1^2u_4$ 13	$7u_3u_4$
$-\frac{115200}{138240}$	$-\frac{1}{29030400} -\frac{1}{4354560}$	$\overline{0} = \frac{1}{87091200} = \frac{1}{207}$	$3600u_1$
$239u_3u_5$ 60	$61u_4^2$ u_9	$17u_4u_5$ $89u_3u_6$	$709u_3^2u_4$
$-\frac{1451520u_1^2}{1451520u_1^2} - \frac{5800}{5800}$	$\frac{1}{6080u_1^2} - \frac{1}{138240u_1^3} - \frac{1}{3}$	$\overline{87072u_1^3} - \overline{3225600u}$	$\frac{3}{1} = \frac{1}{3225600u_1^4}$
$1291u_3u_4^2 = 1001a_4^2$	$u_3^2 u_5 = 197 u_5 u_6 = 163$	$3u_3u_8$ 2069 u_4u_7	$2153u_3^3u_4$
$-\frac{1}{138240u_1^5} - \frac{1}{138240u_1^5}$	$\frac{1}{40u_1^5} - \frac{1}{387072u_1^5} - \frac{1}{967}$	$\frac{1}{680u_1^5} = \frac{1}{5806080u_1^5}$	$-\frac{1}{28800u_1^7}$.
			(3.3.2)

We also computed the genus 5 free energy; see in Appendix A.

For the particular examples of enumerating squares, hexagons, octagons on a genus g surface, one can use (3.3.1), (3.3.2), (A.0.3) to obtain the combinatorial numbers. We checked that these numbers agree with those in [8]. This gives some evidences of validity of the Main Conjecture for g = 3, 4, 5.

Remark 3.3.3 The genus 1, 2, 3 terms of the GUE free energy with even couplings were also derived in [13, 14, 26] for the particular case of only one nonzero coupling (i.e., in the framework of enumeration of 2m-gons). To the best of our knowledge, explicit formulae for higher genus ($g \ge 4$) terms, even in the case of the particular examples, were not available in the literature.

A Explicit formula for F_5

$F_5(u_1,\ldots,u_{13}) =$	
$109514u_2^{12}$	$1352u_2^{11}$ $181628u_3u_2^{10}$ $2593u_2^{10}$ $42691u_3u_2^{9}$
$1215u_1^{16}$	$-\frac{1}{81u_1^{14}} + \frac{1}{405u_1^{15}} - \frac{1}{9450u_1^{12}} + \frac{1}{540u_1^{13}}$
$+\frac{16091u_2^9}{187110u^{10}}-$	$\frac{93460u_4u_2^9}{720u^{14}} + \frac{600763u_3u_2^8}{415800u^{11}} + \frac{134599u_5u_2^8}{4860u^{13}} - \frac{274289u_2^8}{255467520u^8}$
$567100u_1^{-1}$	$1292150u^2u^8$ 201510u u 1^7 20171441u 1^7 151 u^7
$-\frac{567199u_4u_2^2}{12}$	$-\frac{1322159u_3^2u_2^2}{14}+\frac{391519u_3u_4u_2^2}{12}+\frac{2471441u_5u_2^2}{11}-\frac{151u_2^2}{12}$
$24300u_1^{12}$	$1620u_1^{14}$ 972 u_1^{13} 475200 u_1^{11} 399168 u_1^{16}
$999473u_3u_2^7$	$2825u_4u_2^7 1149739u_3^2u_2^7 161353u_6u_2^7 12916717u_3^3u_2^6$
$-2838528u_1^9$	$-\frac{1}{5544u_1^{10}} - \frac{1}{8640u_1^{12}} - \frac{1}{34020u_1^{12}} + \frac{1}{19440u_1^{13}}$
$319877u_3u_2^6$	$31781177u_3u_4u_2^6 + 4549471u_4u_2^6 + 515032871u_5u_2^6$
$+\frac{159667200u_1^7}{159667200u_1^7}$	$+ \frac{475200u_1^{11}}{42577920u_1^8} + \frac{3832012800u_1^9}{3832012800u_1^9}$
$8477461u_7u_2^6$	$263u_2^6 27484783u_6u_2^6 1098923921u_3^2u_2^6$
$+\frac{12830400u_1^{11}}{12830400u_1^{11}}$	$+\frac{1}{23950080u_1^4} - \frac{1}{29937600u_1^{10}} - \frac{1}{425779200u_1^{10}}$
$5713573u_3u_5$	$u_2^6 = 12051881u_4^2u_2^6 + 173831501u_3^3u_2^5 + 6013615u_3^2u_2^5$
$-77760u_1^{12}$	

	$28957u_3u_2^5$ 2	$276125491u_3u_4$	u_2^5 1024	$43u_4u_2^5$	$53761259u_4u_5u_5u_5u_5u_5u_5u_5u_5u_5u_5u_5u_5u_5u$	ι_2^5
	$+\frac{1}{21288960u_1^5}+-$	$182476800u_1^9$	-+ 23950	$00800u_1^6$ +	$3326400u_1^{11}$	
	$102985067u_3u_6u_6u_6u_6u_6u_6u_6u_6u_6u_6u_6u_6u_6u$	$u_2^5 + 84091529a$	$u_7 u_2^5$	u_2^5 1	$11532541u_5u_2^5$	$5117003u_6u_2^5$
	$+$ 9979200 u_1^{11}	+ 63866880	$00u_1^9 + \overline{633}$	$\frac{1}{3600u_1^2} = -$	$479001600u_1^7$	$182476800u_1^8$
	$19847231u_4^2u_2^5$	$2268769u_8u_2^5$	1125727	$7817u_3u_5u_5$	${}^{5}_{2}$ 14531719 u_{3}^{2}	$u_4 u_2^5$
	$2494800u_1^{10}$	$29937600u_1^{10}$	9123	$8400u_1^{10}$	$-36288u_{1}^{1}$	2
	$2322129119u_3^3u_2^4$	$\frac{1}{2}$ 1211 $u_3^2 u_2^4$	8782249	$99u_3u_4^2u_2^4$	$728671781u_3^2u_3^2u_3^2u_3^2u_3^2u_3^2u_3^2u_3^2$	$u_5 u_2^4$
	$+$ 1277337600 u_1^9	$-+\frac{1}{2027520u_1^6}$	+ 1197	$504u_1^{11}$	$12773376u_1^1$	1
	$1046529431u_4u_5$	$_{5}u_{2}^{4}$ 11085336	$611u_3u_6u_2^4$	231617	$u_6 u_2^4 + 1215667$	$u_7 u_2^4 + 1096859 u_9 u_2^4$
	$\phantom{00000000000000000000000000000000000$		$58800u_1^9$	+ 5474304	$40u_1^6 \stackrel{-}{-} 2554675$	$520u_1^7 \stackrel{-}{}^{-} \overline{153280512u_1^9}$
	$29u_2^4$	$47u_3u_2^4$ 29	$783u_4u_2^4$	$24971u_{5}$	$5u_2^4$ 4435112	$51u_3u_4u_2^4$
	$\frac{127733760}{15}$	$\overline{596672u_1^3} - \overline{638}$	$866880u_1^4$	12773376	$50u_1^5 - 191600$	$06400u_1^7$
_	$68301953u_3u_5u_5^2$	$\frac{4}{2}$ _ 5935327 u_8	$\frac{u_2^4}{2} - \frac{13315}{2}$	$50489u_4^2u_2^4$	$5682077u_4u_6$	$5u_2^4$
	$212889600u_1^8$	3832012800	$u_1^8 = 6386$	$568800u_1^8$	$2721600u_1^{10}$	0
_	$-\frac{1095679399u_3^2u_4}{1000000000000000000000000000000000000$	$\frac{1}{4}u_2^4 - \frac{15499108}{15499108}$	$51u_3u_7u_2^4$	48727433	$\frac{377u_5^2u_2^4}{100} - \frac{5475}{100}$	$560589u_3^4u_2^4$
	$19353600u_1^{10}$	136857	$600u_1^{10}$	3832012	$2800u_1^{10}$ 23	$22432u_1^{12}$
_	$+\frac{u_1^2u_2^3}{1}+\frac{6976}{1}$	$\frac{51129u_3u_4^2u_2^3}{2} +$	752920390	$\frac{u_4^2 u_2^3}{1} + \frac{881}{1}$	$17996169u_3^3u_4u_2^3$	
	3991680 6	$967296u_1^9$	287400960	$00u_1^{0}$	$63866880u_1^{11}$	2
_	$+\frac{1049u_4u_2^3}{11055040002}+$	$2479931059u_3^2$	$\frac{u_5u_2^3}{2} + \frac{383}{2}$	$370113u_3u_3$	$\frac{5u_2^3}{6} + \frac{51887531}{6}$	$u_4 u_5 u_2^3$
	$119750400u_{1}^{2}$	319334400 <i>u</i>	u_1^{s} 9	58003200u		800 <i>u</i> ₁
_	$+\frac{2279u_5u_2^3}{19160064u_1^3}+\frac{1}{2}$	$\frac{6316161u_3u_6u_5}{319334400u_1^7}$	$\frac{3}{2} + \frac{255273}{8709}$	$\frac{371u_5u_6u_2^3}{91200u_1^9}$	$+\frac{6169u_6u_2^3}{76640256u_1^4}$	$+ \frac{791123u_4u_7u_2^3}{3870720u_1^9}$
	$369691943u_3u_8u_8u_8u_8u_8u_8u_8u_8u_8u_8u_8u_8u_8u$	$u_2^3 = 284539u_9$	u_2^3 70	$37u_3u_2^3$	$2141u_3^2u_2^3$	$42149u_7u_2^3$
-	$+$ 3832012800 u_1^9	$\frac{1}{2} + \frac{1}{191600640}$	$\frac{1}{0u_1^7} - \frac{1}{2128}$	$\frac{3}{389600u_1}$ -	$-\frac{3}{1520640u_1^4}$ -	$\overline{71850240u_1^5}$
	$1229917u_3u_4u_2^3$	$52781u_8u_2^3$	1086475	$59u_3^3u_2^3$ 2	$2883761u_4u_6u_2^3$	-
-	$638668800u_1^5$	$-\frac{1}{79833600u_1^6}$	47900	$160u_1^7$ –	$8294400u_1^8$	
	$764936639u_3^2u_4u_5$	$u_2^3 = 179669059$	$9u_3u_7u_2^3$	$573019u_1$	${}_{0}u_{2}^{3}$ 4064511	$47u_5^2u_2^3$
	$638668800u_1^8$	9580032	$200u_1^8$ –	104509440	$00u_1^8 - 191600$	$6400u_1^8$
_	$1848263u_4^3u_2^3$	$626921467u_3^2u_4$	$_{6}u_{2}^{3}$ _ 3430	$)962641u_3^4u_3^4u_3^4u_3^4u_3^4u_3^4u_3^4u_3^4$	u_2^3 _ 353380846	$551u_3u_4u_5u_2^3$
	$467775u_1^{10}$	$106444800u_1^1$	0 12	$7733760u_1^{10}$) 191600	$06400u_1^{10}$
_	$+\frac{419193u_3^5u_2^2}{419193u_3^5u_2^2}+-$	$u_1^4 u_2^2 + 23$	$669u_3^2u_2^2$	$+\frac{1783923}{1}$	$\frac{3u_3u_4^2u_2^2}{4u_2^2} + \frac{500}{4u_1^2}$	$47u_4^2u_2^2$
	$14336u_1^{11}$ 6	386688 1277	$7337600u_1^2$	78848	$300u_1^7$ 1064	$44800u_1^4$
_	$+\frac{1238492531u_{3}u_{3}}{12}$	$\frac{2}{5}u_2^2 + \frac{18188001}{18188001}$	$\frac{15u_3^3u_4u_2^2}{15u_3^3u_4u_2^2}$	$+\frac{73978651}{}$	$\frac{1u_3^2u_4u_2^2}{1} + \frac{3337}{1}$	$\frac{u_3u_4u_2^2}{u_3u_4u_2^2}$
	$1277337600u_{1}^{9}$	12773	$3376u_1^9$	638668	$800u_1^6$ 456	$1920u_1^3$
_	$+\frac{5302619u_3^2u_5u_2^2}{200412000}$	$+\frac{15916878976}{12552566}$	$\frac{u_4^2 u_5 u_2^2}{u_5 u_2} + \frac{3}{2}$	$30953u_3u_5u_5u_5u_5u_5u_5u_5u_5u_5u_5u_5u_5u_5u$	$\frac{u_2^2}{4} + \frac{203383903}{12556}$	$u_3 u_4 u_6 u_2^2$
	30412800 <i>u</i> ₁		$00u_1^s$	45619200 <i>u</i>	127733	$\frac{3760u_1^3}{2}$ 2012 7 2
_	$+\frac{29758507u_5u_6u_5}{628668800}$	$\frac{5}{2} + \frac{2758742836}{62966994}$	$\frac{u_{\tilde{3}}u_{7}u_{2}}{00u^{9}} + \frac{2}{10}$	2069131 <i>u</i> ₄ 0	$\frac{u_7 u_2^2}{u_7} + \frac{289 u_3 u_8}{10000}$	$\frac{3u_2^2}{\sqrt{7}} + \frac{60127u_8u_2^2}{0580022000u^4}$
	$038008800u_1^2$	03800880	$500u_1^2$	102	$u_1 = 19008i$	$u_1 = 958003200u_1^*$
_	$+\frac{3313u_9u_2^2}{45610200a^5}+\frac{3}{2}$	$\frac{1143(u_{11}u_{2}^{2})}{248264800.7}$ -	$\frac{5u_1u_3u_2^2}{2128806}$ -	$-\frac{19u_4u_2^2}{54742040}$	$\frac{1}{2} - \frac{1}{2822012800}$	$-\frac{223u_6u_{\tilde{2}}}{0122840a^2}$
	$40019200u_1^2$ 3	040004000 <i>u</i> 1	4140090	04140040	0002012000	u_1 9123840 u_1

$247u_7u_2^2 \qquad 317077u_3u_6u_2^2 \qquad 998201u_3^3u_2^2 \qquad 5091301u_4u_5u_2^2 \qquad 742733u_5^2u_2^2$
$-\frac{11612160u_1^3}{10612160u_1^3} - \frac{1063866880u_1^5}{638668800u_1^5} - \frac{106444800u_1^6}{106444800u_1^6} - \frac{106444800u_1^6}{10644400u_1^6} - \frac{10644400u_1^6}{1064400u_1^6$
$19631u_{10}u_2^2 1295627u_3u_7u_2^2 7277719u_4u_6u_2^2 20664649u_3u_4u_5u_2^2 9854357u_3^2u_6u_2^2$
$-\frac{1}{174182400u_1^6} - \frac{1}{212889600u_1^6} - \frac{1}{638668800u_1^6} - \frac{1}{8870400u_1^8} - \frac{1}{13305600u_1^8} - \frac{1}{13000u_1^8} - \frac{1}{13000u_1^8} - \frac{1}{13000u_1^8} - \frac{1}$
$13316641u_4^3u_2^2 1293697u_5u_7u_2^2 328171u_3u_9u_2^2 45787691u_3^4u_2^2 6093371u_6^2u_2^2$
$- \underbrace{26611200u_1^8}_{26611200u_1^8} - \underbrace{53222400u_1^8}_{53222400u_1^8} - \underbrace{106444800u_1^8}_{106444800u_1^8} - \underbrace{425779200u_1^8}_{425779200u_1^8}$
$18837199u_4u_8u_2^2 \qquad 2912894597u_3^2u_4^2u_2^2 \qquad 1661611993u_3^3u_5u_2^2$
$- 1277337600u_1^8 - 116121600u_1^{10} - 127733760u_1^{10} - 12777300u_1^{10} - 12777300u_1^{10} - 12777300u_1^{10} - 12777300u_1^{10} - 12777300$
$14367497u_3^5u_2 - 3625841u_3^4u_2 - 4721u_3^3u_2 - 274614007u_3u_4^3u_2 - 7368997u_3u_5^2u_2$
$+ \frac{1}{7096320u_1^9} + \frac{1}{127733760u_1^6} + \frac{1}{12773376u_1^3} + \frac{1}{182476800u_1^9} + \frac{1}{70963200u_1^7}$
$927517u_5^2u_2 + 17u_1^3u_3u_2 + 2872733u_3^3u_4u_2 + 20527u_3^2u_4u_2 + 85455011u_4^2u_5u_2$
$+\frac{1}{1916006400u_1^4} + \frac{1}{53222400} + \frac{1}{13305600u_1^7} + \frac{1}{14192640u_1^4} + \frac{1}{638668800u_1^7}$
$+ 1121508611u_3^2u_4u_5u_2 + 159183659u_3^3u_6u_2 + 15535133u_3u_4u_6u_2 + 1516703u_4u_6u_2$
$+ \underbrace{319334400u_1^9}_{110} + \underbrace{212889600u_1^9}_{110} + \underbrace{91238400u_1^7}_{110} + \underbrace{1916006400u_1^4}_{110}$
$2867u_5u_6u_2 + 5833u_6u_2 + 916913u_3^2u_7u_2 + 49661u_3u_7u_2 + 698809u_4u_7u_2$
$+ \frac{1}{1814400u_1^5} + \frac{1}{3832012800} + \frac{1}{19958400u_1^7} + \frac{1}{119750400u_1^4} + \frac{1}{638668800u_1^5}$
$+\frac{82861u_6u_7u_2}{100}+\frac{373u_7u_2}{100}+\frac{650429u_3u_8u_2}{100}+\frac{122413u_5u_8u_2}{100}+\frac{41u_8u_2}{100}$
$45619200u_1^7 91238400u_1 1277337600u_1^5 91238400u_1^7 9580032u_1^2$
$+\frac{38377u_4u_9u_2}{1000}+\frac{3929u_3u_{10}u_2}{1000}+\frac{1091u_{11}u_2}{1000}-\frac{19u_1u_5u_2}{1000}-\frac{23u_1^2u_4u_2}{1000}-\frac{23u_3^2u_2}{1000}$
$53222400u_1^7$ $14515200u_1^7$ $174182400u_1^5$ 13305600 15966720 21288960
$-\frac{4801u_3u_4u_2}{4000} - \frac{5497u_4^2u_2}{4000} - \frac{26833u_3u_5u_2}{40000} - \frac{323u_9u_2}{40000} - \frac{10961u_3u_6u_2}{40000} - \frac{64907u_4u_5u_2}{40000} - \frac{10961u_3u_6u_2}{400000} - \frac{10961u_3u_6u_2}{4000000} - \frac{10961u_3u_6u_2}{400000000} - \frac{10961u_3u_6u_2}{4000000000000000000000000000000000000$
$766402560u_1 63866880u_1^2 212889600u_1^2 68428800u_1^3 79833600u_1^3 273715200u_1^3$
$-\frac{727u_{10}u_2}{10} - \frac{1883363u_3^2u_5u_2}{10} - \frac{3705967u_3u_4^2u_2}{10} - \frac{7u_{12}u_2}{10} - \frac{14227u_6^2u_2}{10} - \frac{16217u_4u_8u_2}{10} - \frac{16217u_4u_8u_8}{10} - \frac{16217u_4u_8u_8}{10} - \frac{16217u_8u_8}{10} - 162$
$116121600u_1^4 159667200u_1^5 239500800u_1^5 4976640u_1^6 7096320u_1^6 7884800u_1^6$
$-\frac{13001u_3u_9u_2}{1000000000000000000000000000000000000$
$15206400u_1^6 \qquad 53222400u_1^6 \qquad 106444800u_1^6 \qquad 159667200u_1^6 \qquad 638668800u_1^6$
$-\frac{1144789u_4u_5^2u_2}{2} - \frac{946477u_3^2u_8u_2}{2} - \frac{176093453u_3^3u_5u_2}{2} - \frac{454075417u_3^2u_4^2u_2}{2} - \frac{28069627u_3u_4u_7u_2}{2} - \frac{176093453u_3^2u_5u_2}{2} - \frac{176093453u_5^2u_5u_2}{2} - \frac{176093453u_5^2u_5u_5u_2}{2} - \frac{176093453u_5^2u_5u_2}{2} - \frac{176093453u_5u_2}{2} - \frac{176093u_5u_2}{2} - \frac{176093u_5u_2}{2} - \frac{176093u_5u_2}{2} - \frac{176093453u_5u_2}{2} - 176093453u_5$
$\frac{11612160u_{1}^{\circ}}{45619200u_{1}^{\circ}} \frac{159667200u_{1}^{\circ}}{212889600u_{1}^{\circ}} \frac{319334400u_{1}^{\circ}}{319334400u_{1}^{\circ}}$
$-\frac{80480347u_3u_5u_6u_2}{200000000000000000000000000000000000$
$638668800u_1^{\circ} 1277337600u_1^{\circ} 1576960u_1^{10} 691200u_1^{\prime} 91238400u_1^{4}$
$+\frac{275599u_3u_4^3}{225222222}+\frac{800453u_4^3}{1012222422}+\frac{571213u_5^3}{22222222222}+\frac{64482661u_3^3u_4^2}{2222222222222}+\frac{1217u_4^2}{1142222222222222}+\frac{22264879u_3u_5^2}{12222222222222}+\frac{1217u_4^2}{122222222222222}+\frac{1217u_4^2}{122222222222222}+\frac{1217u_4^2}{1222222222222222}+\frac{1217u_4^2}{12222222222222222}+\frac{1217u_4^2}{1222222222222222}+\frac{1217u_4^2}{1222222222222222222222}+\frac{1217u_4^2}{12222222222222222222222222}+\frac{1217u_4^2}{122222222222222222222222222222222222$
$3379200u_1^i$ 1916006400 u_1^i 383201280 u_1^i 60825600 u_1^j 1149603840 1277337600 u_1^j
$+\frac{7003u_5^2}{2420264000}+\frac{66653u_3u_6^2}{222}+\frac{u_1^2u_3}{120667200}+\frac{35113699u_3^2u_5}{251156400}+\frac{2932793u_4^2u_5}{12772227000}+\frac{4643u_3u_5}{1016006400}$
$348304800u_1^2 = 28385280u_1^2 = 15900720 = 85155840u_1^2 = 1277337600u_1^2 = 1916000400$
$+\frac{50505149u_3u_4u_5}{150667200u_7}+\frac{16211u_3u_4u_5}{0520022u_4}+\frac{1807u_4u_5}{70822600u_4}+\frac{570989u_3^2u_6}{14102640u_7}+\frac{13u_1^2u_6}{174182400}+\frac{62701u_3^2u_6}{106444800u_4}$
$103001200u_1 \qquad 9000032u_1 \qquad 1300000u_1 \qquad 14132040u_1 \qquad 104162400 \qquad 1004446000u_1^2 \qquad 8660u_4u_4 \qquad 102011u_4u_4u_4 \qquad 20521u_4u_4 \qquad 104291u_4u_4u_4 \qquad 4717u^2u_4$
$+\frac{6009u_3u_6}{628668800_{44}}+\frac{102911u_3u_4u_6}{25481600_{45}}+\frac{20521u_4u_6}{628668800_{42}}+\frac{104281u_4u_5u_6}{14102640_{45}7}+\frac{4717u_5u_7}{6082560_{45}5}$
$050000000u_1 = 55461000u_1^2 = 0580088000u_1^2 = 14192040u_1^2 = 00825000u_1^2$
$+\frac{1559599u_{4}u_{7}}{60825600u_{7}}+\frac{0911u_{1}u_{7}}{11406028400}+\frac{29011u_{3}u_{7}}{1016006400u_{2}^{2}}+\frac{318517u_{3}u_{5}u_{7}}{70822600u_{7}^{2}}+\frac{299u_{6}u_{7}}{1680600u_{7}^{2}}+\frac{1030877u_{3}u_{4}u_{8}}{495770000u_{7}^{2}}$
$00025000u_1^+$ 11490050400 1910000400 u_1^- (9855000 u_1^+ 1089000 u_1^- 425779200 u_1^+

_	$31u_5u_8$	$7697u_3^2u_9$ _ 440	$547u_4u_9$ 7	$1u_3u_{10}$	$13u_{10}$ 1	$13u_{11}$
Т	$\overline{237600u_1^5}$ $^+$	$15206400u_1^7 \stackrel{-}{}^+ 6386$	$668800u_1^5 \stackrel{-}{-} 272$	$21600u_1^5 \stackrel{-}{-} 746$	$49600u_1^2 \stackrel{-}{}^{-} \overline{3483}$	$364800u_1^3$
	u_{13}	$23u_1^2u_3^2$ u	$u_1^4 u_4 = 2011 u_4$	$u_1 u_3 u_4$ 79	$241u^2$	$u_{1}^{3}u_{5}$ u_{1}^{8}
	$29859840u_1^5$	$-\frac{1}{21288960}$ $-\frac{1}{1300}$	636800 - 6386	668800 - 6386	68800 - 958003	$\frac{1}{3200} = \frac{1}{1277337600}$
	$361u_9$	$5707u_{3}^{3}$	$2099u_3^2u_4$	$931u_4u_7$	$13261u_3u_4^2$	$10277u_3^2u_5$
	$547430400u_1$	$-3832012800u_1$	$15966720u_1^2$	$22809600u_1^3$	$54743040u_1^3$	$\overline{58060800u_1^3}$
	$17527u_3u_8$	$113089u_5u_6$	u_{12}	$2533u_3u_9$	$8017u_{6}^{2}$	$8189u_4u_8$
	$\overline{958003200u_1^3}$	$\frac{1916006400u_1^3}{1916006400u_1^3}$	$-\frac{1}{4976640u_1^4}$	$106444800u_1^4$	$\overline{141926400u_1^4}$	$-\frac{141926400u_1^4}{141926400u_1^4}$
	$30553u_5u_7$	$809239u_3^3u_4$	$11u_3u_{11}$	$22349u_4u_5^2$	$701u_5u_9$ 2	$2143u_6u_8$
	$\overline{319334400u_1^4}$	$-\frac{106444800u_1^5}{106444800u_1^5}$	$1741824u_1^6$ - 3	$\frac{1}{3193344u_1^6} - \frac{1}{1}$	$\overline{8247680u_1^6} - \overline{36}$	$5495360u_1^6$
	$775u_4u_{10}$	$520931u_3^3u_5$ 24	$44457u_4^2u_6$ 68	$59977u_3u_4u_7$	$3697u_7^2$	$1138241u_3u_5u_6$
	$\overline{41803776u_1^6}$	$-\frac{1}{42577920u_1^6}-\frac{1}{42}$	$2577920u_1^6$ – 1	$0.06444800u_1^6$	$109486080u_1^6$	$127733760u_1^6$
	$185851u_3^2u_8$	$10135361u_3^2u_4^2$	$336827u_3^2u_4u$	$_{6}$ 3980637 u_{3}^{4}	$u_4 = 653701 u_2$	4
	$127733760u_1^6$	$425779200u_1^6$	$2956800u_1^8$	7884800	$\overline{u_1^8} = \overline{34214400}$	u_1^8
	$877403u_3^3u_7$	$8134913u_3u_4^2u_5$	$22151509u_3^2u$	$\frac{2}{5}$ 4543 u_3^6		$(\Lambda 0.3)$
	$42577920u_1^8$	$45619200u_1^8$	$319334400u_1^8$	$\overline{8192u_1^{10}}$.		(A.0.3)

References

- Bessis, D., Itzykson, C., Zuber, J. B. (1980). Quantum field theory techniques in graphical enumeration. Advances in Applied Mathematics, 1 (2), 109–157.
- [2] Brézin, E., Itzykson, C., Parisi, P., Zuber, J.-B. (1978). Planar diagrams. Commun. Math. Phys., 59, 35–51.
- [3] Buryak, A. (2013). Dubrovin–Zhang hierarchy for the Hodge integrals. Commun. Number Theory and Physics, 9 (2015) 239-271. Preprint arXiv: 1308.5716.
- [4] Carlet, G., Dubrovin, B., Zhang, Y. (2004). The extended Toda hierarchy. Mosc. Math. J., 4 (2), 313–332.
- [5] Dijkgraaf, R., Witten, E. (1990) Mean field theory, topological field theory, and multi-matrix models, Nucl. Phys. B 342 (1990) 486-522.
- [6] Dubrovin, B. (1996). Geometry of 2D topological field theories. In "Integrable Systems and Quantum Groups", Editors: Francaviglia, M., Greco, S. Springer Lecture Notes in Math. 1620, 120– 348.
- [7] Dubrovin, B. (2009). Hamiltonian perturbations of hyperbolic PDEs: from classification results to the properties of solutions. In: New trends in in Mathematical Physics. Selected contributions of the XVth International Congress on Mathematical Physics. Sidoravicius, Vladas (Ed.), Springer Netherlands, 231–276.
- [8] Dubrovin, B., Yang, D. (2016). Generating series for GUE correlators. Preprint arXiv: 1604.07628.

- [9] Dubrovin, B., Yang, D. Remarks on intersection numbers and integrable hierarchies. I. Quasitriviality. to appear.
- [10] Dubrovin, B., Zhang, Y. (2001). Normal forms of hierarchies of integrable PDEs, Frobenius manifolds and Gromov–Witten invariants. Preprint arXiv: math.DG/0108160.
- [11] Dubrovin, B., Zhang, Y. (2004). Virasoro symmetries of the extended Toda hierarchy. Comm. Math. Phys., 250 (1), 161–193.
- [12] Dubrovin, B., Liu, S.-Q., Yang, D., Zhang, Y. (2016). Hodge integrals and tau-symmetric integrable hierarchies of Hamiltonian evolutionary PDEs. Advances in Mathematics, 293, 382–435.
- [13] Ercolani, N. M. (2011). Caustics, counting maps and semi-classical asymptotics. Nonlinearity, 24 (2), 481–526.
- [14] Ercolani, N. M., McLaughlin, K. D. T.-R., Pierce, V. U. (2008). Random matrices, graphical enumeration and the continuum limit of the Toda lattices. Comm. Math. Phys., 278 (1), 31–81.
- [15] Ercolani, N. M., Pierce, V. U. (2012). The continuum limit of Toda lattices for random matrices with odd weights. Comm. Math. Sci. 10, 267–305.
- [16] Faber, C., Pandharipande, R. (2000). Hodge integrals and Gromov-Witten theory. Inventiones mathematicae, 139 (1), 173–199.
- [17] Gopakumar, R., Vafa, C. (1999). On the gauge theory/geometry correspondence, Adv. Theor. Math. Phys., 5, 1415–1443.
- [18] Liu, C. C. M., Liu, K., Zhou, J. (2003). A proof of a conjecture of Mariño–Vafa on Hodge integrals. Journal of Differential Geometry, 65 (2), 289–340.
- [19] Liu, C. C. M., Liu, K., Zhou, J. (2004). On a proof of a conjecture of Mariño–Vafa on Hodge Integrals. Mathematical Research Letters, 11 (2), 259–272.
- [20] Liu, C. C. M., Liu, K., Zhou, J. (2006). Mariño–Vafa formula and Hodge integral identities. Journal of Algebraic Geometry, 15 (2), 379–398.
- [21] 't Hooft, G. (1974). A planar diagram theory for strong interactions. Nucl. Phys. B 72, 461–473.
- [22] 't Hooft, G. (1974). A two-dimensional model for mesons. Nucl. Phys. B 75, 461–470.
- [23] Mehta, M. L. (1991). Random Matrices, 2nd edition, Academic Press, New York.
- [24] Mariño, M., Vafa, C. (2002). Framed knots at large N. Contemporary Mathematics, 310, 185–204.
- [25] Okounkov, A., Pandharipande, R. (2004). Hodge integrals and invariants of the unknot. Geometry & Topology, 8 (2), 675–699.
- [26] Pierce, V. U. (2013). Continuum limits of Toda lattices for map enumeration. In: Algebraic and Geometric Aspects of Integrable Systems and Random Matrices, edited by Anton Dzhamay, Kenichi Maruno, Virgil U. Pierce; Contemporary Mathematics 593, 1–29.
- [27] Witten, E. (1991). Two-dimensional gravity and intersection theory on moduli space. Surveys in Differential Geometry (Cambridge, MA, 1990), (pp. 243–310), Lehigh Univ., Bethlehem, PA.

[28] Zhou, J. (2010). Hodge integrals and integrable hierarchies. Letters in Mathematical Physics, 93 (1), 55–71.

E-mails: dubrovin@sissa.it, dyang@sissa.it